


10th International School of Organometallic Chemistry

SUMMER SCHOOL
Camerino, Italy
05-09 September 2015
<http://d7.unicam.it/isoc/>



University of Camerino
School of Pharmacy
School of Science and Technology

Interdivisional Group of Organometallic Chemistry

Organometallic Chemistry at the Forefront of Societal Challenges



Title Organometallic Chemistry at the forefront of societal challenges
ISBN 9788867680245

Printed by

Tipografia INPRINTING s.r.l. – via Le Mosse, 62032 Camerino (MC)

Tel. 0737 633487 – e-mail: info@inprintingsrl.com

in August 2015

Organometallic Chemistry at the Forefront of Societal Challenges

The ISOC series is the most important school on organometallic chemistry at the European level, organized under the auspices of EuCheMS (the European Association for Chemical and Molecular Sciences) and the Interdivisional Group of Organometallic chemistry of the Italian Chemical Society. The aim is to encourage the presence of young researchers and Ph.D students both from University and Industry, including those not directly involved in organometallic research projects, in order to bring together young researchers and distinguished European scientists as a contribution to the important goal of increasing the transfer of knowledge at a high level between different European countries and different generation of scientists. The 10th edition of ISOC will focus on the role of organometallic chemistry in finding solutions to many of the major societal challenges in the 21st century: from the development of sustainable energy solutions to the mitigation of climate change, from the synthesis of effective therapeutics to the production of new materials ranging from novel polymers to nanomaterials, from the generation of industrial feedstocks to the remediation of the environment. The scientific community is increasingly being stimulated to tackle problems of practical interest and the society will undoubtedly rely on advances made in the field of organometallic chemistry.



TIMETABLE

| 5 th sept | 6 th sept | 7 th sept | 8 th sept | 9 th sept |
|--|---|--|---|---|
| <p>9.00-13.00</p> <p>Registration</p> | <p>9.00-10.30</p> <p>Olivia Reinaud <i>Coordination chemistry in biomimetic cavities</i></p> <p>10.30-11.00</p> <p>Coffee break</p> <p>11.00-12.30</p> <p>Jieping Zhu <i>Palladium-catalyzed domino processes for the synthesis of heterocycles</i></p> <p>12.30</p> <p>Lunch</p> | <p>9.00-10.30</p> <p>Jose Luis Mascareñas <i>New advances in metal catalyzed cycloadditions</i></p> <p>10.30-11.00</p> <p>Coffee break</p> <p>11.00-12.30</p> <p>Nuno Maulide <i>Molecular gymnastics: organic synthesis with rearrangements</i></p> <p>12.30</p> <p>Lunch</p> | <p>8.30-10.00</p> <p>Jacques Maddaluno <i>Organolithium chemistry: an analytical resurrection</i></p> <p>10.00-11.30</p> <p>Axel Jacobi von Wangelin <i>The significance of reductive chemical processes: Examples from industry, research, and future societies</i></p> <p>11.30-11.45</p> <p>Coffee Break</p> <p>11.45-13.15</p> <p>F. Matthias Bickelhaupt <i>Theoretical organometallic chemistry: new concepts and design principles</i></p> <p>13.15</p> <p>Lunch</p> | <p>8.30-10.00</p> <p>Greta R. Patzke <i>Transition metal oxocluster catalysts: from structure-activity relationships to artificial photosynthesis</i></p> <p>10.30-12.00</p> <p>Tobin Jay Marks <i>Interface science of unconventional photovoltaic devices</i></p> |
| <p>15.00-15.30</p> <p>Opening session</p> <p>15.30-17.00</p> <p>Pierre Braunstein <i>Metal-Ligand Interplay for the design of the first and second coordination spheres: fundamental aspects and applications</i></p> <p>17.00-17.30</p> <p>Coffee break</p> <p>17.30-19.00</p> <p>Enzo Alessio <i>The battle against cancer with metal compounds</i></p> <p>20.30</p> <p>Welcome Dinner</p> | <p>14.30-16.00</p> <p>Gianluca Farinola <i>Organometallic approaches to conjugated molecules and polymers for optoelectronics</i></p> <p>16.00-16.30</p> <p>Coffee Break</p> <p>16.30-18.00</p> <p>Gerhard Erker <i>Frustrated Lewis pair chemistry - principles and recent developments</i></p> <p>18.00-19.00</p> <p>Flash Presentation</p> <p>19.00-20.00</p> <p>Poster session</p> <p>20.30</p> <p>Dinner</p> | <p>14.30-16.00</p> <p>Bas de Bruin <i>New catalytic developments involving carbene reactivity</i></p> <p>16.00-16.30</p> <p>Coffee Break</p> <p>16.30-18.00</p> <p>Flash Presentation</p> <p>18.00-20.00</p> <p>Poster session</p> <p>20.30</p> <p>Dinner</p> | <p>15.00</p> <p>Social Tour</p> <p>20.30</p> <p>SOCIAL Dinner</p> | |

TIMETABLE

Saturday, 5 September:

- 15.00: Opening Session
15.30: Prof. Pierre Braunstein “*Metal-Ligand Interplay for the design of the first and second coordination spheres: fundamental aspects and applications*”
17.00: Coffee break
17.30: Prof. Enzo Alessio “*The battle against cancer with metal compounds*”
20.30: Welcome Dinner

Sunday, 6 September:

- 09.00: Prof. Olivia Reinaud “*Coordination chemistry in biomimetic cavities*”
10.30: Coffee break
11.00: Prof. Jieping Zhu “*Palladium-catalyzed domino processes for the synthesis of heterocycles*”
12.30: Lunch
14.30: Prof. Gianluca Farinola “*Organometallic approaches to conjugated molecules and polymers for optoelectronics*”
16.00: Coffee break
16.30: Prof. Gerhard Erker “*Frustrated Lewis pair chemistry - principles and recent developments*”
18.00: Flash Presentation
19.00: Poster session
20.30: Dinner

Monday, 7 September:

- 09.00: Prof. Jose Luis Mascareñas “*New advances in metal catalyzed cycloadditions*”
10.30: Coffee break
11.00: Prof. Nuno Maulide “*Molecular gymnastics: organic synthesis with rearrangements*”
12.30: Lunch
14.30: Prof. Bas de Bruin “*New catalytic developments involving carbene reactivity*”
16.00: Coffee break
16.30: Flash Presentation
18.00: Poster session
20.30: Dinner

Tuesday, 8 September:

- 08.30: Prof. Jacques Maddaluno “*Organolithium chemistry: an analytical resurrection*”
10.00: Prof. Axel Jacobi von Wangelin “*The significance of reductive chemical processes: Examples from industry, research, and future societies*”
11.30: Coffee break
11.45: Prof. F. Matthias Bickelhaupt “*Theoretical organometallic chemistry: new concepts and design principles*”
13.15: Lunch
15.00: Social Tour
20.30: Social dinner

Wednesday, 9 September:

- 08.30: Prof. Greta R. Patzke “*Transition metal oxocluster catalysts: from structure-activity relationships to artificial photosynthesis*”
10.30: Prof. Tobin Jay Marks “*Interface science of unconventional photovoltaic devices*”
12.00: Prizes and Closing Ceremony

ORGANIZING COMMITTEE



Prof. Augusto Cingolani
Honorary President of ISOC



Prof. Claudio Pettinari
Chair



Prof. Fabio Marchetti
Co-chair



Prof. Riccardo Pettinari



Prof. Adriano Pizzabocca



Prof. Roberto Ballini




Prof. Marino Petrini



Prof. Enrico Marcantoni



Prof. Corrado Di Nicola



Dr. Jessica Palmucci



Dr. Francesca Condello



Dr. Agnese Petrini



Dr. Nello Mosca

Advisory Board

Pierre Braunstein - University of Strasbourg, France
Ernesto Carmona - University of Sevilla, Spain
Jan Čermák - Institute of Chemical Process Fundamentals - Czech Republic
Antonella Dalla Cort - University La Sapienza, Roma, Italy
Kees Elsevier - University of Amsterdam, Netherlands
Roberto Gobetto - University of Torino, Italy
Helena Grennberg - Uppsala University, Sweden
David Milstein - The Weizmann Institute of Science, Israel
Luis A Oro - University of Zaragoza, Spain
Jun Okuda - RWTH Aachen University, Germany
Robin A. Perutz - University of York, United Kingdom
Maurizio Peruzzini - ICCOM CNR Firenze, Italy
Claudio Pettinari - University of Camerino, Italy
Rinaldo Poli - LCC, CNRS, Toulouse, France
Armando Pombeiro - University of Lisbon, Portugal
Michelangelo Scalone - F. Hoffmann - La Roche AG
Valerio Zanotti - University of Bologna, Italy

Scientific Committee

Prof. Marino Basato
Prof. Silvia Bordoni
Prof. Antonella Dalla Cort
Prof. Emanuela Licandro
Prof. Alceo Macchioni
Dr. Alessandro Mordini
Prof. Roberto Gobetto
Prof. Fabio Ragaini
Prof. Claudio Pettinari
Prof. Enrico Marcantoni

SPEAKERS



Prof. Pierre Braunstein France, CNRS-Université de Strasbourg

Metal-Ligand Interplay for the design of the first and second coordination spheres: fundamental aspects and applications




Prof. Enzo Alessio Italy, University of Trieste

The battle against cancer with metal compounds



Prof. Olivia Reinaud France, University of Paris Decartes

Coordination chemistry in biomimetic cavities



ÉCOLE POLYTECHNIQUE
FÉDÉRALE DE LAUSANNE

Prof. Jieping Zhu Switzerland, Institute of Chemical Sciences and Engineering, École Polytechnique Fédérale de Lausanne

Palladium-catalyzed domino processes for the synthesis of heterocycles



Prof. Gianluca Farinola Italy, University of Bari

Organometallic approaches to conjugated molecules and polymers for optoelectronics

SPEAKERS



Prof. Gerhard Erker



Germany, University of Münster

Frustrated Lewis pair chemistry - principles and recent developments



Prof. Jose Luis Mascareñas



Spain, CIQUS University of Santiago de Compostela

New advances in metal catalyzed cycloadditions



Prof. Nuno Maulide




Austria, University of Vienna

Molecular gymnastics: organic synthesis with rearrangements



Prof. Bas de Bruin



Netherlands, University of Amsterdam

New catalytic developments involving carbene reactivity



Prof. Jacques Maddaluno



France, CNRS-University of Rouen



Organolithium chemistry: an analytical resurrection

SPEAKERS



Prof. Axel Jacobi von Wangelin Germany, University of Regensburg

The significance of reductive chemical processes: Examples from industry, research, and future societies



Prof. F. Matthias Bickelhaupt Netherlands, VU University Amsterdam and Radboud University Nijmegen

Theoretical organometallic chemistry: new concepts and design principles



Prof. Greta R. Patzke Switzerland, University of Zurich

Transition metal oxocluster catalysts: from structure-activity relationships to artificial photosynthesis



NORTHWESTERN UNIVERSITY
USA, Northwestern University

Interface science of unconventional photovoltaic devices

LECTURES


Metal-Ligand Interplay for the Design of the First and Second Coordination Spheres: Fundamental Aspects and Applications

Prof. Pierre Braunstein


^a Laboratoire de Chimie de Coordination, Institut de Chimie (UMR 7177 CNRS),
Université de Strasbourg, 4 rue Blaise Pascal, 67081 Strasbourg, France
Email: braunstein@unistra.fr

Global energy needs In addition to the stronger metal-ligand interactions of the covalent- or dative-type that characterize the first coordination sphere of a metal centre, weaker interactions, such as H-bonding and metallophilic interactions, can occur that can often be viewed as shaping the second coordination sphere of the metal. Although the energies involved in the latter case are smaller, the influence of these weaker interactions on the chemical and/or physical properties of a metal complex has long been recognized.

We shall focus on the properties of transition metal complexes that are directly associated with the presence of functional ligands able to chelate a metal centre through at least two chemically-different donors. This has direct implications on the reactivity of the other ligands, see *e.g.* positions A and B)



Based on the 'phosphine mimicry' of N-heterocyclic carbenes (NHCs), a number of functional NHCs have been investigated and their rich chemistry is attracting increasing attention (see **I**). Chelating or bridging modes can be envisaged, depending on the spacer used. NHC-based « pincer-type » ligands also fall into this category (see **II**) (for simplicity, all the donor groups in **I** and **II** have been taken as neutral, 2 e donors, but this does not have to be the case).



We will examine specific features associated with different donor functions connected to the NHC core. These will include oxygen-, sulfur-, imine-, and phosphine-based functionalities.

Some of these ligands can be tailored to behave as short-bite ligands and this will give us the opportunity to discuss unique features associated with the presence of an additional functionality in short-bite diphosphine ligands.

The Battle Against Cancer and Other Diseases with Metal Compounds

Prof. Enzo Alessio

Department of Chemical and Pharmaceutical Sciences, University of Trieste, Via Giorgieri 1, 34127 Trieste, Italy.
Email: alessi@units.it

The general aim of the lecture is that of showing the huge potentialities that coordination and organometallic compounds have in the field of medicinal inorganic chemistry. Given the time limit, the lecture will be restricted to therapeutic applications of metal compounds (real or potential), excluding radioactive metals for radiotherapy, and will not deal with diagnostic applications, i.e. the use of metal compounds as imaging agents.

After a general introduction, the first part of the lecture will focus on the most successful inorganic drugs so far, i.e. on Pt anticancer compounds. Their serendipitous discovery, development and, above all, their chemical features and mechanism of action will be treated. The problems and limitations of Pt anticancer drugs will lead to introduce anticancer compounds based on other metals and, in particular, on ruthenium, both coordination complexes (e.g. KP1019 and NAMI-A) and organometallic species (e.g. piano-stool compounds). At the end of this part of the lecture, a general classification of metal based drugs depending on their mode of action will be proposed.

From this point onward the lecture will focus on organometallic species exclusively, and the distinctive features of organometallic compounds that make them particularly appealing for being investigated in medicinal inorganic chemistry will be highlighted. Different approaches to the development of organometallic drugs – not limited to the anticancer field – will be shown, in particular the bio-isosteric replacement of phenyl rings with metallocene fragments in bioactive molecules (such as in the antimalarial drug *ferroquine* and in the anticancer compounds named *ferrocifens*). In addition to potential anticancer species, examples of organometallic compounds investigated as selective enzyme inhibitors and as antibacterial drugs will be shown.

In the final part of the lecture, the use of light for the activation of metal compounds in biological systems (i.e. photodissociation of ligands, caged complexes) will be addressed.

For each argument treated, the lecture will provide updated references.

LECTURES

Coordination Chemistry in biomimetic cavities

Prof. Olivia Reinaud


Laboratoire de Chimie et Biochimie Toxicologiques et Pharmacologiques, CNRS UMR 8601, Université Paris Descartes, Sorbonne Paris Cité,
45 rue des Saints Pères, 75006 Paris.

Email: Olivia.Reinaud@parisdescartes.fr

Group of Supramolecular Bioinorganic Chemistry <http://www.biomedicale.univ-paris5.fr/umr8601/-Chimie-Bioinorganique-.html>

Supramolecular chemistry concerns the reversible assembly of discrete entities through the establishment of multiple weak interactions between the different components. As these phenomena are fundamental in the biological world, Nature has been a major source of inspiration for chemists involved in the supramolecular field. Biomimetic inorganic chemistry is mainly focused on mimicking the first sphere coordination environment of the metal ion. Little information is available concerning the influence, or even the control, that the microenvironment provided by a protein can have on the reactivity of the metal. Exploration of this aspect through the construction of cavity-based complexes will be presented and discussed.¹ Hence, the concepts of “funnel” complex and “bowl” complex will be illustrated by selected examples that gave rise to:

- The elaboration of receptors that are highly selective for neutral molecules² and efficient even in the highly competitive medium, which is water,³
- A new proposed mechanism for copper mono-oxygenases,⁴
- The design of systems allowing the control of hetero-multinuclear binding of metal ions,⁵
- A biomimetic strategy for the selective functionalization of a cavity: Guest Covalent
- Capture by a Host.⁶



References

- [1] *Supramolecular Bioinorganic Chemistry*, J.-N. Rebilly and O. Reinaud in *Supramolecular Chemistry: From Molecules to Nanomaterials* (SMC174), Wiley, 2012.
- [2] D. Coquière, S. Le Gac, U. Darbost, O. Sénéque, I. Jabin, O. Reinaud, *Org. Biomol. Chem.*, **2009**, *7*, 2485-2500.
- [3] O. Bistri and O. Reinaud, *Org. Biomol. Chem.*, **2015**, *13*, 2849-2865
- [4] N. Le Poul, Y. Le Mest, I. Jabin and O. Reinaud, *Acc. Chem. Res.*, **2015**, DOI: 10.1021/acs.accounts.5b00152.
- [5] J.-N. Rebilly, B. Colasson, O. Bistri, D. Over, O. Reinaud, *Chem. Soc. Rev.* **2015**, *44*, 467-489; J.-N. Rebilly, O. Reinaud, *Supramol. Chem.* **2014**, 454-479.
- [6] B. Colasson and O. Reinaud, *J. Am. Chem. Soc.* **2008**, *130*, 15226-15227.

Palladium-catalyzed Domino Processes for the Synthesis of Heterocycles: Case of Indoles and Indole Alkaloids

Prof. Jieping Zhu

Laboratory of Synthesis and Natural Products, Institute of Chemical Sciences and Engineering,
Ecole Polytechnique Fédérale de Lausanne, EPFL-SB-ISIC-LSPN, CH-1015 Lausanne, Switzerland.

Email: jieping.zhu@epfl.ch

The indole nucleus is a ubiquitous heterocycle found in many bioactive natural products, pharmaceuticals and agrochemicals. The synthesis and functionalization of indoles have attracted chemists for over a century and remain an active research area.¹ This presentation will be divided into two parts. The first part will focus on the Pd(II)-catalyzed *bis-nucleophilic* addition across a triple bond. The concept led to the development of efficient synthesis of indoloisoquinolones, 3-alkynylindoles and cyclizative cross-coupling reactions under aerobic conditions.² Detailed mechanistic and kinetic studies allowed us to propose subsequently a robust and low-catalyst loading process for the access to this important class of heterocycles. The second part will discuss the development of a Pd-catalyzed decarboxylative cross coupling of aliphatic carboxylic acids with vinyltriflates for the synthesis of 1-aryl-1-vinyl-alkanes and the application of this methodology in the total synthesis of monoterpene indole alkaloids.³

References:

- [1] Cacchi, S.; Fabrizi, G. *Chem. Rev.* **2011**, *111*, PR215-PR283.
- [2] Yao, B.; Wang, Q.; Zhu, J. *Angew. Chem. Int. Ed.* **2012**, *51*, 5170-5174. Yao, B.; Wang, Q.; Zhu, J. *Angew. Chem. Int. Ed.* **2012**, *51*, 12315-12315; Yao, B.; Wang, Q.; Zhu, J. *Angew. Chem. Int. Ed.* **2013**, *52*, 12992-12996. Yao, B.; Wang, Q.; Zhu, J. *Chem. Eur. J.* **2014**, *20*, 12255-12261. Ha, T. M.; Bo, Yao, Wang, Q.; Zhu, J. *Org. Lett.* **2015**, *17*, 1750-1753.
- [3] Xu, Z.; Wang, Q.; Zhu, J. *Angew. Chem. Int. Ed.* **2013**, *52*, 3272-3276. Xu, Z.; Wang, Q.; Zhu, J. *J. Am. Chem. Soc.* **2013**, *135*, 19127-19130. Wagnières O.; Xu, Z.; Wang, Q. *J. Am. Chem. Soc.* **2014**, *136*, 15102-15108. Xu, Z.; Wang, Q.; Zhu, J. *J. Am. Chem. Soc.* **2015**, *137*, 6712-6724.

LECTURES

Organometallic Approaches to Conjugated Molecules and Polymers for Optoelectronics

Prof. Gianluca M. Farinola

Università degli Studi di Bari "Aldo Moro", Dipartimento di Chimica, Via Orabona 4, 70126 Bari, Italy.
Email: gianlucamaría.farinola@uniba.it

Progress of organic optoelectronics is largely based on the possibility to optimize the performances of molecular and polymeric organic semiconductors. Not only optical and electronic properties, but also processability and stability can be finely tuned and enhanced by molecular design. On the other hand, industrial production of plastic solar cells,¹ organic light emitting diodes² and other organic electronics devices³ requires synthetic methods scalable up to thousands of kilograms, keeping in consideration costs and environmental impact.

Synthetic protocols of molecular and polymeric semiconductors are commonly based on organometallic reactions which are efficient, versatile and highly selective.⁴

The lecture will discuss the most important organometallic approaches to conjugated oligomers and polymers for applications in organic photonics and electronics. Classical Pd-catalyzed cross-coupling processes (*e.g.* Stille, Cassar-Heck-Sonogashira, Suzuki-Miyaura reactions) as well as more recent methods (*e.g.* direct arylation by C-H bond activation) will be presented for the synthesis of several classes of conjugated polymers and small molecules for optoelectronics.

Besides discussing scope and limitations of the organometallic methods, the lecture will introduce the issue of industrial scalability of the synthetic processes,¹ an important aspect that has been often neglected in the literature, showing possible approaches to quantify the synthetic accessibility⁵ as a key parameter in the evaluation of the real practical significance of the organometallic protocols.

References

- [1] Marzano, G.; Ciasca, C. V.; Babudri, F.; Bianchi, G.; Pellegrino A.; Po, R.; Farinola G. M. *Eur. J. Org. Chem.* **2014**, *30*, 6583.
- [2] Farinola, G. M.; Ragni, R. *Chem. Soc. Rev.* **2011**, *40*, 3467.
- [3] Operamolla, A.; Farinola, G. M. *Eur. J. Org. Chem.*, **2011**, 423.
- [4] Babudri, F.; Farinola, G. M.; Naso F. *J. Mater. Chem.* **2004**, *14*, 11.
- [5] Osedach, T. P.; Andrew, T. L.; Bulovic, V. *Energy Environ. Sci.*, **2013**, *6*, 711. Po, R.; Bianchi, G.; Carbonera, C.; Pellegrino, A. *Macromolecules*, **2015**, *48*, 453.

Frustrated Lewis Pair Chemistry - Principles and Recent Developments

Prof. Gerhard Erker

Organisch-Chemisches Institut, Universität Münster, Germany Corrensstr. 40, Münster.
Email: erker@uni-muenster.de

Lewis acids and bases usually undergo a neutralizing adduct formation when brought together. However, Lewis acids and bases that feature very bulky substituents at their central atoms can evade this neutralization reaction. This may lead to situations of combinations of active Lewis acids and bases in solution. Such "frustrated Lewis pairs" (FLPs) offer the chance to observe cooperative reactions with added substrates. This has indeed been observed. A variety of phosphane/borane or amine/borane FLPs were shown to undergo unusual reactions with added small molecules. Several examples will be presented including heterolytic dihydrogen cleavage and metal-free catalytic hydrogenation, but also a number of new reactions such as CO reduction by [B]-H boranes or the development of a phospho-Stork chemistry. Eventually some developments toward active metal containing FLPs will be presented and discussed. These use Group 4 metallocenes as Lewis acids, but there are also systems made up by combinations of Group 4 metallocenes and boranes. The utilization of such systems in catalysis will be discussed.

References

- [1] Review: "Frustrated Lewis Pair Chemistry: Development and Perspectives", D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.* **2015**, *54*, 6400-6441, doi: 10.1002/anie.201409800.

LECTURES


New Advances in Metal Catalyzed Cycloadditions

Prof. José Luis Mascareñas

Departamento de Química Orgánica and Centro Singular de Investigación en Química Biolóxica e Materiais Moleculares (CIQUS) C/Jenaro de la Fuente s/n. Campus Vida Universidade de Santiago 15782 Santiago de Compostela, Spain.

Email: josehlmascarenas@usc.es

Cycloaddition reactions allow the direct assembly of cyclic skeletons from acyclic precursors, and therefore represent invaluable tools to streamline the synthesis of complex polycyclic products. While classical cycloadditions are somewhat restricted to specific, electronically complementary reactants, the potential of cycloaddition chemistry can be enormously expanded by taking advantage of the coordination and activation capabilities of metal complexes.¹ In recent years we have unveiled several new cycloaddition modes of alkylidenecyclopropanes by using palladium or nickel catalysts.² We have also developed several platinum or gold-catalyzed reactions of allenes consisting of formal (4 + 3) and (4 + 2) cycloadditions.³ The intermolecular version of these reactions could be better accomplished by using allenamides as cycloaddition partners. These substrates work as efficient two carbon components in a variety of gold-catalyzed cycloadditions with different types of alkenyl derivatives (see scheme).⁴ Most recently, our work in cycloaddition chemistry is moving towards the discovery of "anomalous" annulations involving C-H activation processes.⁵



References

- [1] M. Lautens, W. Klute, W. Tam *Chem. Rev.* **1996**, *96*, 49-92.
- [2] (a) M. Gulías, F. López, J.L. Mascareñas *Pure Appl. Chem.* **2011**, *83*, 495-506. (b) J.L. Mascareñas, M. Gulías, F. López in *Comprehensive Organic Synthesis* 2nd edition, Eds: G. A. Molander, P. Knochel, Vol 5, Oxford: Elsevier; 2014, 595-655.
- [3] F. López, J. L. Mascareñas *Chem. Soc. Rev.* **2014**, *43*, 2904-2915.
- [4] (a) Faustino, H.; López, F.; Castedo, L.; Mascareñas, J. L. *Chem. Sci.* **2011**, *2*, 633-637; (b) J. Francos, F. Grande-Carmona, H. Faustino, J. Iglesias-Sigüenza, E. Díez, I. Alonso, R. Fernández, J. M. Lassaletta, F. López, J. L. Mascareñas *J. Am. Chem. Soc.* **2012**, *134*, 14322-14325; (c) Faustino, H.; Alonso, I.; Mascareñas, J. L.; Lopez, F. *Angew. Chem. Int. Ed.* **2013**, *52*, 6526-6530.
- [5] A. Seoane, N. Casanova, N. Quiñones, J.L. Mascareñas, M. Gulías, *J. Am. Chem. Soc.* **2014**, *136*, 834-837.

Molecular Gymnastics: C-C Bond Formation with Rearrangements

Prof. Nuno Maulide

University of Vienna, Institute of Organic Chemistry, Währinger Strasse 38, 1090 Vienna, Austria.

Email: nuno.maulide@univie.ac.at

The turn of the century brought about a pressing need for new, efficient and clean strategies for the chemical synthesis of biorelevant compounds. Our group has studied the use of various molecular rearrangements and atom-economical transformations as particularly appealing means towards the streamlined synthesis of complex small molecule targets.^{1,2,3}

In this lecture, we will present an overview of our research in these areas and how they provide efficient solutions for total synthesis as well as platforms for the discovery of unusual reactivity.

References

- [1] (a) Luparia, M.; Oliveira, M.T.; Audisio, D.; Frébault, F.; Maulide, N. *Angew. Chem. Int. Ed.* **2011**, *50*, 12631. (b) Audisio, D.; Luparia, M.; Oliveira, M.T.; Frébault, F.; Klütt, D.; Maulide, N. *Angew. Chem. Int. Ed.* **2012**, *51*, 7314. (c) Misale, A.; Niyomchon, S.; Luparia, M.; Maulide, N. *Angew. Chem. Int. Ed.* **2014**, *53*, 7068.
- [2] Huang, X.; Maulide, N. *J. Am. Chem. Soc.* **2013**, *135*, 7312.
- [3] Jurberg, I.D.; Peng, B.; Wüstefeld, E.; Wasserloos, M.; Maulide, N. *Angew. Chem. Int. Ed.* **2012**, *51*, 1950.


LECTURES

New Catalytic Developments Involving Carbene Reactivity

Prof. Bas de Bruin

Van 't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam, P.O. Box 94720 1090 GE Amsterdam, The Netherlands.
Email: b.debruin@uva.nl

This presentation starts with an overview of catalytic reactions involving Fischer- and Schrock-type carbenes. These reactions play an important role in several processes ranging from synthesis of fine chemicals to polymeric materials.



The second part of this presentation focusses on new developments in our own group using both Fischer-type carbenes and a new carbene type: carbene radicals. These new reactions give access to novel reaction patterns that are not only interesting from a fundamental perspective, but also give access to completely new synthetic protocols to synthesize pharmaceuticals and polymers.

Organolithium Chemistry: an Analytical Resurrection

Prof. Jacques Maddaluno

CNRS, University & INSA Rouen
e-mail : jmaddalu@crihan.fr

Organolithium reagents are now commonplace in modern organic synthesis. In fact, their popularity has progressively pushed these reagents in the far back of the “routine tools” drawer, to a point where they are barely considered worthy of an active contemporary research in organometallic chemistry.

We wish to show in this lecture that indeed the synthetic applications of organolithium, in particular as deprotonating agents, has been largely explored and mastered in the last 25 years. But we also want to emphasize that relatively little is known about the structure of these reagents in solution while this parameter is likely to exert a significant influence on their reactivity. Since modern analytical and theoretical tools have opened refined accesses to the intimate organization of these species in solution and to some aspects of their reactivity, we are convinced it is now time for an analytical resurrection of this class of compounds.

In the first part of this lecture, we will remind the participants with the fundamentals of this type of chemistry, in terms of stability and reaction conditions. We will then examine in relative details the main families of deprotonation reactions by organolithium bases (and the factors influencing the chemoselectivity, the ortho-lithiation, lateral lithiation) and their mechanisms. We will next consider another important class of reactions: the halogen-lithium exchange and its main characteristics.

In the second part of the lecture, we will spend the remaining time on more exploratory aspects of the chemistry of organolithium reagents, based on our own researches. In particular, we will give a rapid overview on the carbolithiation of alkynes and the use of chiral mixed aggregates for the enantioselective addition of organolithium reagents on carbonyl compounds. These results make an extensive use of modern analytical tools, in particular NMR ones, as well as of theoretical (DFT) computations to provide with a wide description of this domain of organometallic chemistry.



LECTURES


The Significance of Reductive Chemical Processes: Examples from Industry, Research & Future Societies

Prof. Axel Jacobi von Wangelin

Institute of Organic Chemistry, University of Regensburg Universitaetsstrasse 31, 93040 Regensburg (Germany).


Email: axel.jacobi@ur.de

The aerobic conditions on our planet have enabled the accumulation of oxidized matter whereas reduced chemicals constitute most valuable energy carriers. The future shortages of energy-rich resources make efficient reductive conversions of available raw materials into valuable chemicals and fuels one of the greatest challenges of modern societies.




Over the past decades, **metal-catalyzed reductions and related reductive processes** have emerged as key synthetic methods for the preparation of diverse building blocks, fine chemicals, pharmaceuticals, agrochemicals and materials. Such reactions are mostly operated with noble or toxic catalyst metals (Pd, Ni, Cu, Co, Rh, Pt etc.) - which is only partially due to their good catalytic activity. The relatively high stability of noble metals in lower oxidation states has allowed facile catalyst syntheses, implementations of structural variations, and spectroscopic and mechanistic studies that led to a deeper understanding of the underlying elemental steps and consequently to a steady progress in the development of more effective catalyst systems and reaction parameters. However, the commonly employed catalysts exhibit detrimental effects on the environment and generate high operational costs. **Iron catalysts** are a promising yet under-utilized alternative for reductive transformations. From an abstract thermodynamic perspective it is obvious that base metals (e.g. Fe) exhibit a much stronger reductive power in low oxidation states than commonly employed noble metals.

Capitalize on reductive power:



.. and sustainability:



Fe accounts for 5% of the earth's crust, it's the 10th most abundant element of the universe and has no critical hazard profile. The Fe-catalyzed N₂ fixation (Haber-Bosch) and CO reduction (Fischer-Tropsch) are the two largest technical reductions. But only very recently, other efficient Fe-catalyzed reductions, hydrogenations, and reductive coupling reactions have been developed.

Theoretical Organometallic Chemistry: New Concepts and Design Principles

Prof. F. M. Bickelhaupt

Amsterdam Center for Multiscale Modeling (ACMM), VU University Amsterdam, and Institute for Molecules and Materials (IMM), Radboud

University Nijmegen.

Email: f.m.bickelhaupt@vu.nl

The aim of this work is to understand¹ the factors that determine the activity and selectivity of transition-metal catalysts in oxidative-addition steps that occur in bond activation and cross-coupling reactions. We studied the effect of varying the metal M along the d¹⁰ metals of groups 9, 10 and 11, in combination with varying the number as well as the type of ligands, using relativistic density functional theory (DFT)² and the activation strain model (ASM).¹




Figure 1. Schematic illustration of concepts presented in this lecture.

Three concepts for tuning a catalyst's activity emerge: (i) d-regime catalysts; (ii) s-regime catalysts; and (iii) bite-angle flexibility (see Figure 1, left). These concepts explain why the effect of one and the same ligand on a catalyst's activity can be completely different, even opposite, for catalysts from different electronic regimes. Furthermore, I will point out why not the bite angle itself, but its flexibility is decisive for the catalyst's activity. Finally, I show how the latter is influenced not only by steric repulsion but also by what we designate "steric attraction" (see Figure 1, right).

References

- [1] (a) L. P. Wolters, F. M. Bickelhaupt, *WIREs Comput. Mol. Sci.* **2015**, *5*, online (*open access*). (b) I. Fernandez, F. M. Bickelhaupt, *Chem. Soc. Rev.* **2014**, *43*, 4953-4967; (c) F. M. Bickelhaupt, *J. Comput. Chem.* **1999**, *20*, 114-128.
- [2] (a) G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. Fonseca Guerra, S. J. A. van Gisbergen, J. G. Snijders, T. Ziegler, *J. Comput. Chem.* **2001**, *22* 931-967; (b) F. M. Bickelhaupt, E. J. Baerends, In: *Reviews in Computational Chemistry*; K. B. Lipkowitz and D. B. Boyd, Eds.; Wiley-VCH: New York, **2000**, Vol. 15, pp. 1-86; (c) C. Fonseca Guerra, J.-W. Handgraaf, E. J. Baerends, F. M. Bickelhaupt, *J. Comput. Chem.* **2004**, *25*, 189-210.

Transition Metal Oxocluster Catalysts: From Structure-Activity Relationships to Artificial Photosynthesis

Prof. Greta R. Patzke

University of Zurich, Institute of Inorganic Chemistry, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland.
Email: greta.patzke@aci.uzh.ch

Inspired **Scope:** Artificial photosynthesis has emerged as a promising and elegant strategy to convert solar energy resources into clean and sustainable fuels.^[1] After decades of research into the mechanisms of water splitting processes, water oxidation still remains a challenging bottleneck towards technological implementations. This renders the design of robust, economic and efficient water oxidation catalysts a highly relevant current research target.

Background: The introduction is followed by an overview of recent progress in artificial photosynthesis setups. The pros and cons of heterogeneous and molecular catalysts for water splitting are briefly compared. Next, up-to-date concepts for the construction and application of molecular water oxidation and reduction catalysts (WOCs and WRCs) are discussed. The oxygen-evolving complex (OEC) of photosystem II is introduced as a leading design paradigm for the construction of bio-inspired WOCs.^[2]

Polyoxometalates as WOCs and WRCs: This section sets out with a brief introduction to the structural and analytical chemistry of polyoxometalates (POMs) as a key family of multi-functional oxoclusters which attract increasing interest for catalytic applications.^[3] Next, latest developments on POM-based WOCs are summarized and their interplay between hetero- and homogeneous catalysis is discussed. Two case studies^[4] on the structure-activity relationships (SAR) of POM catalysts round this section off and shed light on the challenging interaction of POMs with standard photosensitizers, along with the key influence of subtle factors, such as crystallographic disorder, on the water oxidation performance. Furthermore, the upcoming use of POMs as WRCs is highlighted, followed by a systematic investigation of structural features of small polyoxotungstates and their effects on the photocatalytic water reduction activity.

Molecular cubanes as bio-inspired WOCs: The final part summarizes the present efforts to translate the structural features of the OEC into molecular cubanes as platforms to investigate the parameters and mechanisms of bio-related artificial photosynthesis. Given that the construction of active WOCs based on direct OEC replicas remains a major challenge, considerable current activities are directed on $\{\text{Co}_4\text{O}_4\}$ -based cubane WOCs. Latest results on the SAR of $\{\text{Co(III)}_4\text{O}_4\}$ -WOCs and their stability under operational conditions are discussed. Moreover, the catalytic performance of a new $\{\text{Co(II)}_4\text{O}_4\}$ -WOC with a flexible OEC-inspired ligand architecture is introduced in more detail as a model system for translating bio-mimetic features into stable WOCs. Solution stability tests, activity parameters, and ligand exchange as well as flash photolysis studies provide an insight into the mechanisms associated with Co-cubane WOCs. Future trends for molecular oxocluster WOC design are presented in the outlook.

References

- [1] (a) S. Berardi, S. Drouet, L. Francàs, C. Gimbert-Surinach, M. Guttentag, C. Richmond, T. Stoll, A. Llobet, *Chem. Soc. Rev.* **2014**, 43,7501; (b) Z. Yu, F. Li, L. Sun, *Energy Env. Sci.* **2015**, 8, 760.
[2] (a) J.-R. Shen, *Annu. Rev. Plant Biol.* **2015**, 66, 23; (b) C. Zhang, C. Chen, H. Dong, J.-R. Shen, H. Dau, J. Zhao, *Science* **2015**, 348, 690.
[3] A. Sartorel, M. Bonchio, S. Campagna, F. Scandola, *Chem. Soc. Rev.* **2013**, 42, 2262.
[4] (a) F. Evangelisti, P.-E. Car, O. Blacque, G. R. Patzke, *Catal. Sci. Technol.* **2013**, 12, 3117; (b) K. von Allmen, R. Moré, R. Müller, J. Soriano-López, A. Linden, G. R. Patzke, *ChemPlusChem* **2015**, DOI: 10.1002/cplu.201500074.
[5] F. Evangelisti, R. Güttinger, R. Moré, S. Lubber, G. R. Patzke, *J. Am. Chem. Soc.* **2013**, 135, 18734.

Interface Science of Soft Matter Photovoltaics

Prof. Tobin J. Marks

Department of Chemistry, Materials Research Center, and the Argonne-Northwestern Solar Research Center
Northwestern University, Evanston IL 60208, USA

The ability to fabricate molecularly tailored interfaces with nanoscale precision offers means to selectively modulate charge transport, molecular assembly, and exciton dynamics at hard matter-soft matter and soft-soft matter interfaces. Such interfaces can facilitate transport of the “correct charges” while blocking transport of the “incorrect charges” at the electrode-active layer interfaces of organic photovoltaic cells. This interfacial tailoring can also suppress carrier-trapping defect densities at interfaces and stabilize them with respect to physical/thermal de-cohesion. For soft matter-soft matter interfaces, interfacial tailoring can also facilitate exciton scission and photocurrent generation in such cells. In this lecture, challenges and opportunities in organic photovoltaic interface science are illustrated for four specific and interrelated areas of research: 1) controlling charge transport across hard matter(electrode)-soft matter interfaces in organic photovoltaic cells, 2) controlling charge transport by specific active layer nano/microstructural organization in the bulk active material and at the electrodes, 3) controlling exciton dynamics and carrier generation at donor-acceptor interfaces in the active layer, 4) designing transparent conducting oxide electrodes with improved properties. It will be seen that such rational interface engineering along with improved bulk-heterojunction polymer structures guided by theoretical/computational analysis affords exceptional fill factors, solar power conversion efficiencies greater than 10%, and enhanced cell durability.

Poster 1

Green Valorization of Glycerol by Metal Catalysts

Massimiliano Lupacchini,^{a*} Nicola d'Alessandro,^a Lucia Tonucci^b

^a Department of Engineering and Geology (INGEO), "G.d'Annunzio" University of Chieti-Pescara, Viale Pindaro, 42, 65127 Pescara, Italy.

Email: massimiliano.lupacchini@studenti.unich.it

^b Department of Philosophical, Educational and Economic Sciences, "G.d'Annunzio" University of Chieti-Pescara, Via dei Vestini 31, 66100 Chieti, Italy.


The increasing of biodiesel production leads to a growing supply of the by-product, glycerol. This prompts researchers to investigate for selective chemical transformations of glycerol, aiming to obtain derivatives with high added value and wide-ranging uses as new fuels, polymers and fine chemicals.¹

Several catalytic pathways were proposed to achieve efficient, cheap and "green" deoxygenation, such as selective thermal processing, hydrogenolysis, acid-catalyzed dehydration and bio-mediated reactions.

However a very competitive methodology, namely, deoxydehydrogenation reaction (DODH) has been only marginally investigated. DODH can remove two adjacent hydroxyl groups from vicinal diols to afford the corresponding unsaturated derivatives which, in the case of glycerol, is the allyl alcohol, a useful building block commonly employed for many applications.

In the present study we describe a green conversion of glycerol in allyl alcohol using several rhenium-based catalysts: ReCl₅, ReI₅, [(C₆H₅)₃P]₂ReO₂I.

Recovery of the reaction products was achieved using an ice-cooled water trap collecting the gas flow coming from the reaction vessel, where air was continuously bubbled (Figure 1).²



Scheme 1. Schematic representation of the reactor employed for rhenium catalyzed DODH of glycerol.

References

- [1] (a) Posada, J. A.; Rincón, L. E.; Cardona, C. A. *Bioresour. Technol.* **2012**, *111*, 282; (b) Long, Y. D.; Fang, Z. *Biofuels, Bioprod. Biorefin.* **2012**, *6*, 686; (c) Clacens, J. M.; Pouilloux, Y.; Barrault, J. *Appl. Catal.* **2002**, *227*, 181; (d) Martin, A.; Richter, M. *Eur. J. Lipid Sci. Technol.* **2011**, *113*, 100; (e) Nomanbhay, A.; Hussain, R.; Rahman, M. M.; Palanisamy, K. *Adv. Nat. Appl. Sci.* **2012**, *6*, 802.
[2] Canale, V.; Tonucci, L.; Bressan, M.; D'Alessandro, N. *Catal. Sci. Technol.* **2014**, *4*, 3697–3704.

Poster 2

The Palladium-Catalyzed Heck arylation of Homoallylic Substrates in the Presence of Ionic Liquids Containing (-)-Menthyl Group

A. Wirwis,^{a*} A.M.Trzeciak,^a J. Feder-Kubis^b


^a Faculty of Chemistry, University of Wrocław, 14 F. Joliot – Curie St., 50-383 Wrocław, Poland.

Email: anna.wirwis@chem.uni.wroc.pl

^b Faculty of Chemistry, Wrocław University of Technology, Wybrzeże Wyspiańskiego 27, 50-370 Wrocław, Poland

Palladium-catalyzed Heck reaction has been well known as one of the most important methods for the formation of new C-C bonds.¹ It is interesting to use allylic alcohols as substrates in Heck reaction, because it is an easy way to obtain carbonyl compounds which are useful intermediates for the synthesis of pharmaceuticals, fragrances or natural products.²

The first Heck reaction products, arylated allylic alcohols, can undergo isomerization to carbonyl compounds, for example to 4-aryl-2-butanones.^{3,4}



In this paper we present new method of synthesis of saturated arylated ketones, involving a cross-coupling between homoallylic alcohols and aryl halides (Fig.1). We used PdCl₂(COD) as a catalyst and chiral ionic liquids containing (-)-menthyl group as co-catalyst. The reaction was carried out at 80°C and 24h in the presence of weak base – NaHCO₃. We tested a lot of variation of parameters such as solvents or time in order to obtain high yield of desired product.

For comparison, an alternative pathway leading to saturated carbonyl compound under the same conditions was studied. Using unsaturated ketones instead of homoallylic alcohols it was possible to obtain unsaturated ketones which were next hydrogenated in the same catalytic system. In both reactions we observed positive effect of the presence of ionic liquids.

References

- [1] Beletskaya, P.I.; Cherpakov, V.A.; *Chem.Rev.* **2000**, *100*, 3009.
[2] Liu, S.; Thomson, N.; Pettman, A.; Hyder, Z.; Mo, J.; Xiao, J.; *J.Mol.Catal. A: Chemical* **2008**, *279*, 210.
[3] Alacid, E.; Najera, C.; *Adv.Synth.Catal.* **2007**, *349*, 2572.
[4] Viviano, M.; Glasnov, N.T.; Reichart, B.; Tekautz, G.; Kappe, O.C.; *Am.Chem.Soc.* **2011**, *15*, 858.

Poster 3

Divalent Lanthanides and Alkaline-Earth Metal Complexes in Intermolecular Hydrophosphination Reactions of Olefins

I.V. Basalov,^{a,*} D.M. Lyubov,^a G.K. Fukin,^a A.V. Cherkasov,^a S.-C. Roşca,^b Y. Sarazin,^b J.-F. Carpentier^b and A.A. Trifonov^a


^a G. A. Razuvaev Institute of Organometallic Chemistry of Russian Academy of Sciences, Nizhny Novgorod, Russia.

Email: basalov.vania@yandex.ru


^bUniversity of Rennes, France.

Hydrophosphination reactions it is an addition of the P-H bond of the phosphorous compounds to C-C unsaturated bond, which is a convenient synthetic method of forming a C-P bond. Such phosphorus-organic compounds find a wide range of applications, from ligands in metal complexes to building blocks in medical chemistry and catalysts of organic synthesis.

We obtained and structurally characterized a series of stable heteroleptic amide complexes of divalent ytterbium, samarium, and calcium containing aminoethyl-phenolate ligand systems.



The resulting compounds exhibit high catalytic activity, productivity and selectivity in the intermolecular hydrophosphination reaction between styrene and phenylphosphine with high TON and TOF, up to 0.04% catalyst loading.



Acknowledgements - This work was supported by RFFI (12-03-93109-NtsNIL.a) grant of the President of the Russian Federation for the state support of young Russian scientists (MK-5702.2015.3).

References

[1] I. V. Basalov, V. Dorcet, G. K. Fukin, J.-F. Carpentier, Y. Sarazin, A. A. Trifonov, *Chem. Eur. J.* **2015**, *21*, 6033 – 6036.

Poster 4

Complexes of Yttrium Supported by Bulky Amido-Imino and en-Diamido Ligands. Synthesis, Structure, Reactivity and Catalytic Activity in Intermolecular Hydrophosphination and Hydroamination Reactions of Olefins

Alexander A. Kissel,^{a,*} Tatyana V. Mahrova,^a Dmitry M. Lyubov,^a Anton V. Cherkasov,^a Georgy K. Fukin,^a Alexander A. Trifonov,^a Iker Del Rosal,^b Laurent Maron^b

^a G.A. Razuvaev Institute of Organometallic Chemistry of Russian Academy of Sciences, 603950 Nizhny Novgorod, Russia.

Email: okishvegan@mail.ru

^b Université de Toulouse, INSA, 31077 Toulouse, France.

Due to the high biological and industrial importance of nitrogen-containing and phosphorus-containing molecules, of great interest to the scientific and industrial research had development of effective methods for the synthesis of these molecules. Hydroelementation reactions, that is the addition of E-H (E = N, P) functionalities across C-C multiple bonds, are atom-economic processes for the synthesis of valuable compounds that may be one of the prospective application fields of organo-rare-earth complexes.

The compounds containing bond Ln-C and Ln-H are of interest, both in terms of basic research and the use of these complexes in the conversion reaction of unsaturated substrates, due to the high reactivity of these derivatives of lanthanides.

A monoanionic amido-imino and dianionic en-diamido ligand systems were successfully employed for the synthesis of alkyl $[(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3)\text{NC}(\text{Me})_2\text{Y}(\text{CH}_2\text{SiMe}_3)(\text{THF})_2]$ (1), $\{[2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{NC}(\text{Me})\text{-C}(\text{Me})\text{N-C}_6\text{H}_5\text{-}i\text{Pr}_2\text{-}2,6]\text{Y}(\text{CH}_2\text{SiMe}_3)_2\}^-\{\text{Li}(\text{THF})_4\}^+$ (2), $[(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3)\text{N}=\text{C}(\text{Me})\text{-C}(\text{CH}_2\text{N}(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3)]\text{Y}(\text{CH}_2\text{SiMe}_3)_2$ (THF) (4) and hydride $\{[(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3)\text{NC}(\text{Me})_2\text{YH}(\text{THF})_2](\text{THF})\}$ (3) complexes (Figure 1).




Figure 1.

Alkyl and hydrido complexes **1**, **2**, **3** and **4** proved to be an efficient catalysts of hydroelementation reactions. They enable hydrophosphination of para-substituted styrenes and 4-vinylpyridine with PhPH₂ and Ph₂PH under mild conditions with formation of the anti-Markovnikov product. Furthermore these complexes initiate hydroamination of para-substituted styrenes with pyrrolidine.

Acknowledgements – This work was supported by the Russian Foundation for Basic Research (Grant no. 12-03-93109-НЦННЦ). DML thanks for a Grant of President of Russian Federation for young scientists (grant no. MK-5702.2015.3).

Poster 5

Metal Complexes with Di(N-Heterocyclic Carbene) Ligands Bearing an *Ortho*-, *Meta*- or *Para*-Phenylene Bridge

Marco Monticelli,^{a,*} Cristina Tubaro,^a Claudia Graiff,^b Gianluca Accorsi^c

^a Dipartimento di Scienze Chimiche, Università degli Studi di Padova, via Marzolo 1, 35131 Padova, Italy.

Email: marco.monticelli.1@studenti.unipd.it

^b Dipartimento di Chimica, Università degli Studi di Parma, Viale delle Scienze 17/A, 43100 Parma, Italy.

^c CNR NANOTEC - Istituto di Nanotecnologia, Polo di Nanotecnologia c/o Campus Ecotekne, via Monteroni - 73100 Lecce, Italy.

Di(N-heterocyclic carbene) ligands (diNHC) are one of the most interesting classes of compounds studied for the syntheses of transition metal complexes with a wide range of applications, spanning from photophysics, catalysis and medicine.¹ Generally the employed diNHCs have a flexible alkyl chain as bridging group between the carbene moieties. In this work, we present our recent results on diNHC ligands with rigid *ortho*-, *meta*- and *para*-phenylene bridging group, which could be interesting because the different geometry of functionalization of the aromatic ring should influence the bridging vs. chelating coordination of the dicarbene.^{2,3} Furthermore, a limited number of examples are present in the literature with these ligands and in most cases they involve the *meta*-phenylene bridge, which can act also as *CCC*-pincer ligand as a consequence of the metalation of the phenylene bridge in the C2 position (with Ru(II), Ir(III) or Pt(II) centers).³ We have synthesized the dinuclear bridging diNHC silver(I) complexes and studied their ability to transfer the diNHC ligands to other metal centers (like Ru(II), Ir(III), Cu(I) and Au(I)). As expected, Ag(I), Au(I) and Cu(I) complexes present the classical linear coordination; the diNHC transmetalation to the Ir(III) and Ru(II) centers is successfully only with the *ortho*-phenylene ligand and affords the chelate complexes (Figure 1). Finally, the solid state structures and the luminescence properties of the Ag(I) and Au(I) complexes will be discussed and compared.




Figure 1.

References

- [1] Biffis, A.; Baron, M.; Tubaro, C. *Adv. Organomet. Chem.* doi:10.1016/bs.adomc.2015.02.002.
- [2] Rit, A.; Pape, T.; Hepp, A.; Hahn, F. E. *Organometallics* **2011**, *30*, 334.
- [3] Raynal, M.; Pattacini, R.; Cazin, C. S. J.; Vallée, C.; Olivier-Bourbigou, H.; Braunstein, P. *Organometallics* **2009**, *28*, 4028.

Poster 6


Mn(III) Complexes Containing Hydrophilic Chiral Ligands

Maria E. Cucciolito, Giovanni Di Mauro, Stefania Esposito, Marco Palmese,* Francesco Ruffo

Dipartimento di Scienze Chimiche, Università di Napoli "Federico II", Italy and Consorzio Interuniversitario di Reattività Chimica e Catalisi (CIRCC), Italy.

Complexes of Mn(III) containing chiral derivatives of salicylethylenediamine (salen) have remarkable stereo-orienting properties in enantioselective catalysis, for example in the asymmetric epoxidation of unsaturated substrates.¹

The environmental impact of these transformations can be reduced by using a well tolerated reaction solvent, such as water, which offers numerous advantages for its cost and safety. On these grounds, the present study aims to prepare a library of Mn(III) complexes (**1**) containing the structural motifs useful for the extension of these enantioselective processes in aqueous environments.³ More precisely, the tetradentate ligand will be based on the chiral frame resulting from *trans*-1,2-cyclohexanediamine or stilbendiamine.



Hydrophilic substituents R will be introduced in the key positions 3 and 3', for prompting both enantioselectivity¹ and water solubility (an example is **1a**).

The compounds will then be evaluated in the asymmetric epoxidation reaction of styrenes in water.

References

- [1] For a "review", see: Katsuki, T. J. *Mol. Catal. A: Chemical*, **1996**, *113*, 87-107.
- [2] Cornils, B.; Herrmann, W. A. (Eds.), *Aqueous-Phase Organometallic Catalysis*, Wiley-VCH, **1998**.
- [3] For a similar approach, see: Sun, Y.; Tang, N. J. *Mol. Catal. A: Chemical*, **2006**, *255*, 171-179.

Poster 7

Biomimetic Association of a Copper Ion and Organic Redox Units Using Calix[6]Arene Based Ligands

Sarah Richard,^{a*} Jean-Noël Rebilly,^a Gaëtan Le Duc,^a Andrea Brugnara,^a Marina El Gohary,^a Benoit Colasson,^a Jean-Noël Rebilly^a and O. Renaud^a

^aLaboratoire de Chimie et Biochimie Pharmacologiques et Toxicologiques, CNRS UMR 8601, Université Paris Descartes, 45 rue des Saints Pères, Paris 75006, France.
Email: sarah.richard@etu.parisdescartes.fr

Calix[6]arenes are interesting platforms that can be functionalized in order to design ligands that mimic the active site of metallo-enzymes.¹ Previously, a calix[6]arene covalently linked to a TMPA unit was used to model the active site of mono-copper oxygenases.² This work aims to develop models of copper oxidases such as galactose oxidase and amine oxidase that associate an organic redox co-factor next to the copper center (a tyrosine residue and a topaquinone, respectively). The strategy presented relies on the synthesis of calix[6]TMPA derivatives that incorporate phenol or quinone units in the hosting cone (Figure 1).




Figure 1

First insights into calix-phenol reactivity highlight the formation of a radical species (Figure 2). The synthesis of the ligands, the host-guest and redox properties of the corresponding copper complexes will be presented.




Figure 2

References

- [1] Rebilly, J.-N.; Renaud, O. *Supramol. Chem.* **2014**, *26*, 454-479.
[2] Thiabaud, G.; Guillemot, G.; Schmitz-Afonso, I.; Colasson, B.; Renaud, O. *Angew. Chem. Int. Ed.* **2009**, *48*, 7383-7386.

Poster 8

Luminescent Metal Tetrazolate Complexes: From Design To Applications

Valentina Fiorini,^{a*} Stefano Stagni,^a Massimiliano Massi^b

^a Department of Industrial Chemistry "Toso Montanari", Viale del Risorgimento 4, 40136 Bologna (BO), Italy.
Email: valentina.fiorini5@unibo.it

^b Department of Chemistry, Curtin University, 6102 Bentley WA, Australia.

In homogeneous The preparation and the study of the physical and chemical properties of coordination compounds containing 5-aryl tetrazolate [R-CN₄⁻] ligands is the focus of our research program. In particular we have demonstrated how these synthetically versatile nitrogen-rich ligands can actually rule the luminescent output of the corresponding Ru(II), Ir(III), Re(I) and Pt(II) – based complexes.^{1,2} The synthetic versatility of tetrazolate derivatives combined with the peculiar sensitivity to electrophiles that is displayed by tetrazolate complexes constitute the basis for the further development of their coordination chemistry and their applicative scenarios. Following a similar approach, we have prepared a new family of anionic Ir(III) tetrazolate complexes and new Re(I)-based species containing ditopic tetrazolate ligands. (Figure 1) The photophysical properties of the new molecules will be presented herein, and particular attention will be dedicated to their use as components for soft salts and to the study of their chemosensing abilities.




Figure 1

References

- [1] Werrett, M. V.; et al. *Inorg. Chem.*, **2014**, *53*, 229.
[2] Werrett, M. V.; et al. *Dalton Trans.*, **2015**, *44*, 8379.

Poster 9

Electro-Synthesis Towards High Energy Intermediates

Colet te Grotenhuis,^{a*} Bas de Bruin^a

^a Homogeneous, Supramolecular and Bio-Inspired Catalysis, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands.

Email: C.teGrotenhuis@uva.nl




Figure 1.

Many (multistep) reactions involve the synthesis of high energy intermediates. This puts a high demand on the environment through energy consumption and waste formation. A possible solution is the use of electro-synthesis. In this case the energy isn't stored in molecules but in electrons. The exact amount of energy can be applied. One possible application can be found in the synthesis of Ritalin. The high energy intermediate shown in Figure 1 can be reached through double deprotonation and oxidation.

Poster 10

New Luminescent Ir(III) Complexes with 1,2,3-Triazolylenes as NHC-Type Ligand

Elia Matteucci,^{a*} Andrea Baschieri,^a Filippo Monti,^b Andrea Mazzanti,^a Nicola Armaroli^b and Letizia Sambri^a

^aDepartment of Industrial Chemistry "Toso Montanari", University of Bologna, Italy.

Email: elia.matteucci2@unibo.it

^bIstituto per la Sintesi Organica e la Fotoreattività, CNR, Bologna, Italy

Triazolylidene derivatives have found a wide range of application, especially in organometallic chemistry and catalysis, due principally to the relatively high covalent contribution to the M–NHC bond and to their strong donating abilities. The success of this class of ligand can be also attributed to easily preparation of the triazole precursor via the readily accessible Cu(I) catalysed 'click' cycloaddition of an alkyne and azide (CuAAC).¹ Subsequent alkylation of the triazole with a suitable alkylating agent is generally selective and yields the 1,3,4-substituted triazolium salt. Metal complexation is readily available too via direct metallation with Ag₂O and subsequent transmetallation with a wide range of metal precursors, giving a whole library of metal-triazolylidene complexes.²

Only few examples of triazolylidene metal complexes that show photochemical properties have been reported.³

Here we report the first examples of luminescent Ir(III) complexes bearing two triazolylidene ligands.




Figure 1.

References


- [1] (a) Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem. Int. Ed.*, **2002**, *41*, 2596; (b) Himo, F.; Lovell, T.; Hilgraf, R.; Rostovtsev, V. V.; Noodleman, L.; Sharpless, K. B.; Fokin, V. V. *J. Am. Chem. Soc.*, **2005**, *127*, 210.
- [2] Donnelly, K. F.; Petronilho, A.; Albrecht, M. *Chem. Commun.*, **2013**, *49*, 1145.
- [3] (a) Schulze, B.; Escudero, D.; Friebe, C.; Siebert, R.; Görls, H.; Köhn, U.; Altuntas, E.; Baumgaertel, A.; Hager, M. D.; Winter, A.; Dietzek, B.; Popp, J.; Gonzalez, L.; Schubert, U. S. *Chem. Eur. J.*, **2011**, *17*, 5494; (b) Leigh, V.; Ghattas, W.; Lalrempuia, R.; Pryce, M.; Müller-Bunz, H.; Albrecht, M. *Inorg.Chem.* **2013**, *52*, 5395.

Functionalization of Ferrite Nanoparticles by Phenanthroline/Palladium Complexes and Their Use as Catalysts in Pharmaceutical Syntheses


Lilian M.Silva Ansaloni,* Francesco Ferretti, Fabio Ragaini

Università degli Studi di Milano, Dipartimento di Chimica, via Golgi 19, Milano
e-mail: lilian.silva@unimi.it

We In the last decade magnetic nanoparticles (MNP) have been widely used as a support for the immobilization of homogeneous catalysts. The small size of these particles allows their thorough dispersion in the reaction media thus nearing the homogeneous environment of soluble catalysts. At the same time, their magnetic properties allow to easily recover the catalyst at the end of the reaction. In our group, we developed a method with high reproducibility for the synthesis of ferrite MNP functionalized with a phosphonic acid bearing a halogen in the ω position to allow further modification. Owing to our interest in carbonylation and reductive cyclization reaction of nitro compounds, catalyzed by palladium/phenanthroline complexes,¹⁻³ we functionalized the protecting layer with a phenanthroline ligand.



$\text{Pd}(\text{OAc})_2$ was then immobilized on the $\text{MNP}(\text{N}^*\text{N})$ and the catalyst tested in the cyclization of nitrostyrenes to indoles using CO as a reductant.⁴ The indole skeleton is present in several molecules of pharmaceutical interesting, one example is fluvastatin.



Acknowledgement: CAPES-BRAZIL Grant BEX-5963-13-3

References

- [1] F. Ferretti, E. Gallo, F. Ragaini, *J. Organomet. Chem.* **771**, (2014), 59.
- [2] F. Ferretti, E. Gallo, F. Ragaini, *ChemCatChem*, (2014), 10.1002/cctc.201500452.
- [3] F. Ragaini, A. Rapetti, E. Visentin, M. Monzani, A. Caselli, S. Cenini, *J. Org. Chem.* **71**, (2006), 71, 3748.
- [4] I. W. Davies, J. H. Smitrovich, R. Sidler, C. Qu, V. Gresham, C. Bazaral, *Tetrahedron* **61**, (2005), 6425.

Metal Bound *Cis* Olefin of Olive Oil Fatty Acid in $[\text{PtI}_2(\text{Me}_2\text{phen})(\eta^2\text{-alkene})]$ Complexes: a Potential Platinum Delivery to Hydrophobic Targets

Claudio Carrara,* Andrea Pizzi, Silvia Sonzini, Emanuela Licandro Sandra A. De Pascali, Laura Del Coco, Chiara R. Girelli*, Paride Papadia, Francesco P. Fanizzi

University of Salento, Dipartimento di Scienze e Tecnologie Biologiche ed Ambientali, – 73100 Lecce – Italy.
Email: chiara.girelli@unisalento.it

It is known that distorted square planar complex of formula $[\text{PtX}_2(\text{Me}_2\text{phen})]$ (Me_2phen , 2,9-dimethyl-1,10-phenanthroline, X = Cl, Br, or I) can react reversibly with olefins giving the corresponding very stable pentacoordinate species of the type $[\text{PtX}_2(\text{Me}_2\text{phen})(\eta^2\text{-olefin})]$.¹ This kind of complexes has a trigonal bipyramidal geometry, with the two halogens in axial positions, and the neocuproine and the olefin bound to platinum in the trigonal plane² (Figure 1a). The reactivity of the square planar species towards olefin is strongly favoured by halogen sterical hindrance which determines distortion of the coordination geometry, increasing in the order $\text{Cl} < \text{Br} < \text{I}$.^{1,3} We studied the reaction between the very sterically hindered square planar species, $[\text{PtI}_2(\text{Me}_2\text{phen})]$, and unsaturated moieties of esterified fatty acid chains naturally occurring in olive oils. We performed experiments directly with extra virgin olive oils (EVOOs) and different standards of methyl esters of fatty acids, as *cis*- Δ^9 methyl oleate and *cis*- Δ^9, Δ^{12} methyl linoleate.




Figure 1. (a) Structure of general pentacoordinate complexes; b) structure of synthesized $[\text{PtI}_2(\text{Me}_2\text{phen})(\eta^2\text{-Methyloleate})]$

The pentacoordinated Pt(II) species (Figure 1b) obtained by the coordination of the $[\text{PtI}_2(\text{Me}_2\text{phen})]$ complex with the *cis* olefin groups of the esterified fatty acid moieties were characterized by 1D and 2D NMR spectroscopy. ¹H, ¹³C, ¹⁹⁵Pt chemical shifts signals were analysed and discussed, suggesting the formation of mono and bis-adducts. Moreover, Diffusion Ordered Spectroscopy (DOSY) NMR experiments were performed to discriminate the different species obtained. Being the potential pharmacological activity of water-soluble pentacoordinate complexes already demonstrated⁴, this preliminary study could suggest a possible use of these pentacoordinate species as prodrugs also on hydrophobic targets.

References

- [1] Fanizzi, F.P.; Intini, F.P.; Maresca, L.; Natile, G.; Lanfranchi, M.; Tiripicchio, A., *J. Chem. Soc. Dalton Trans.* **1991**, 1007.
- [2] Fanizzi, F.P.; Maresca, L.; Natile, G.; Lanfranchi, M.; Tiripicchio, A.; Pacchioni, G. *J. Chem. Soc., Chem. Commun.* **1992**, 4, 333.
- [3] Benedetti, M., Papadia, P.; Girelli, C.R.; De Castro, F.; Capitelli, F.; Fanizzi, F.P. *Inorg. Chim. Acta* **2015**, 428, 8.
- [4] De Pascali, S.A.; Migoni, D.; Papadia, P.; Muscella, A.; Marsigliante, S.; Ciccarese, A.; Fanizzi, F.P. *Dalton Trans.* **2007**, 5077.

Pd Mediated Cross Coupling Reactions for the Synthesis of New MOF Linkers

Davide Balestri,^{a*} Paolo Pelagatti^a

^a Dipartimento di Chimica, Università degli Studi di Parma, Parco Area delle Scienze, 17/A 43124 Parma, Italy.

Email: davide.balestri@studenti.unipr.it

MOFs (Metal Organic Frameworks) are a large class of ultraporous crystalline polymers, generated from the self-assembly of metallic ions or clusters and bridging organic linkers. The wide choice both in organic ligands and metallic nodes leads to an extraordinary modulation of the final structural properties. For these reasons, the design of the organic linkers has a key role in obtaining the desired topology of the MOF network. Pd-catalyzed cross coupling reactions are certainly a powerful tool for the ligand synthesis and provide several advantages over conventional organic synthesis, such as functional group tolerance, one step procedures and, consequently, reduction of wastes. Buchwald and coworkers proposed an easy synthetic pathway for the metal-catalyzed aminations of aryl halides,^{1,2} process that proved to be quite difficult in other reaction conditions. Exploiting this procedure is also possible to form C-N bonds by using very poor nucleophiles, such as amides, combined with aryl chlorides with excellent yields.³ In this communication we report on the successful application of the Buchwald protocol for the construction of novel organic linkers useful for the fabrication of new MOFs. A scheme summarizing the isolated ligands is reported in Figure 1. A series of ditopic or tritopic ligands characterized by different degrees of flexibility, linearity and denticity were isolated and fully characterized. Their use as organic linkers for the fabrication of new MOFs frameworks is currently under investigation in our laboratory.




Figure 1.

References

- [1] D. S. Surry and S. L. Buchwald, *Chem. Sci.*, **2011**, 2, 27–50.
- [2] B. P. Fors, D. A. Watson, M. R. Biscoe, and S. L. Buchwald, *J. Am. Chem. Soc.*, **2008**, 130, 13552–13554.
- [3] B. P. Fors, P. Krattiger, E. Strieter, and S. L. Buchwald, *Org. Lett.*, **2008** Vol. 10, No. 16, 3505–3508.


Manipulating Radicals: Using Cobalt to Steer Radical Reactions

Andrei Chirila,^{a*} Bas de Bruin^a

^a Department of Homogeneous and Supramolecular Catalysis, Van 't Hoff Institute for Molecular Sciences, Faculty of Science, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands.

Email: a.chirila@uva.nl

Cobalt(II) Schiff base complexes have proven to be effective catalysts for radical-type transformations, such as cyclopropanation or aziridination.^{1,2} This work focuses on finding suitable catalysts for the activation of diazo compounds, which can subsequently lead to formation of β -lactams, 2,3-dihydrobenzofuranes, amides, cyclopropanes, etc. This research shows that cobalt(II) catalysts based on ligands such as tetraazaannulenes or salens proved very active towards the activation of diazo-compounds, in some cases outclassing their porphyrin equivalents.³



References

- [1] W.I. Dziki, X. Xu, X.P. Zhang, J.N.H. Reek and B. de Bruin, *J. Am. Chem. Soc.* **2010**, 132, 10891.
- [2] D. Intriери, A. Caselli, and E. Gallo, *Eur. J. Inorg. Chem.* **2011**, 5071.
- [3] N.D. Paul, A. Chirila, H. Lu, X.P. Zhang, B. de Bruin, *Chem. Eur. J.* **2013**, 19, 12953.

DFT Investigation on [FeFe]-Hydrogenases and Biomimetic Models: Implications on the Activation of H₂

F. Arrigoni,* L. Bertini, L. De Gioia, G. Filippi, G. Zampella

Department of Biotechnology and Biosciences, University of Milano-Bicocca, Italy.

[FeFe]-hydrogenases are enzymes that catalyze the reversible oxidation of H₂ with extremely high turnovers.¹ The chance of carrying out the same transformation in a controlled and productive way, has attracted a broad scientific community, since H₂ is a clean and high density fuel. The enzyme active site consists of a di-iron cluster, bound to a Fe₂S₄ cubane. Several biomimetic catalysts have been synthesized, but their activity is much lower than that of the natural system, particularly in the oxidation direction. Remarkably, it was recently showed that the presence of a metallic redox-active ligand has important implications in catalysis, since H₂ activation can be driven by partial spin delocalization among the metallic centers.² Here we present a DFT study on H₂ activation, in both natural and biomimetic systems, in order to highlight the effect of ligands nature on catalysis.

Aminocarbonylation in Aqueous Medium Catalyzed by Palladium(II) Complexes with Ar,Ar'-BIAN Ligand (Ar,Ar'-BIAN = Bis(aryl)acenaphthenequinonediimine)


Przemysław Wójcik,^{a*} Vera Rosar,^b Anna M. Trzeciak,^a Barbara Milani^b

^a Faculty of Chemistry, University of Wrocław, 14 F. Joliot-Curie Str., 50-383 Wrocław, Poland.

Email: przemyslaw.wojcik@chem.uni.wroc.pl


^b Dipartimento di Scienze Chimiche e Farmaceutiche, Università di Trieste, Via Licio Giorgieri 1, 34127 Trieste, Italy.

In the last decades the aminocarbonylation of primary and secondary amines with aryl halides is of significant interest in academia and fine chemical industry (Scheme 1). Products of these reactions – amides – constitute an interesting structural motif which is frequently present in pharmaceuticals (atorvastatin, mosapride),¹ in biologically active compounds having antagonistic action to many hormones^{2,3} and in components for the manufacture of synthetic fibers (nylon, perlon).



Scheme 1. The aminocarbonylation reaction.

Up to now, catalytic activity of palladium complexes incorporating Ar,Ar'-BIAN ligand in aminocarbonylation of aniline with aryl iodides have been never studied. In this work the study of the coordination properties of (3,5-(CH₃)₂C₆H₃)₂-BIAN and (2,6-(CH₃)₂C₆H₃)₂-BIAN ligands towards [PdCl₂(cod)] precursor has been performed. Two complexes: [(3,5-(CH₃)₂C₆H₃)₂-BIAN]PdCl₂ and [(2,6-(CH₃)₂C₆H₃)₂-BIAN]PdCl₂ with Ar,Ar'-BIAN ligand connected to palladium atom in a bidentate mode have been obtained (Scheme 2) and characterized by spectroscopic methods. The catalytic activity of these complexes was tested in aminocarbonylation reaction of aniline with iodobenzene. The results show that, under mild reaction conditions (CO (balloon pressure), 4h, 90 °C) and in green solvent (water), these complexes generate relatively high-yielding and selective catalysts for aminocarbonylation reaction. Analysis of post-catalytic solution by Transmission Electron Microscopy showed the presence of palladium nanoparticles which are generated *in situ* during reaction.



Scheme 2. Palladium catalyst with Ar,Ar'-BIAN ligand.

References

- [1] Lubitz W. et al, *Chem. Rev.* **2007**, 107, 4331–65.
[2] Rauchfuss T.B. et al, *Nat. Chem.* **2011**, 4, 26–30.

References

- [1] A. Brennführer, H. Neumann, M. Beller, *Angew. Chem. Int. Ed.* **2009**, 48, 4114–4133.
[2] T. F. Walsh, R. B. Toupence, J. R. Young, a S. X. Huang, *Bioorg. Med. Chem. Lett.* **2000**, 10, 443–447.
[3] A. Gangjee, Y. Zeng, J. J. McGuire, R. L. Kisliuk, *J. Med. Chem.* **2002**, 45, 1942–1948.

The Tethered Aminohydroxylation of Glycals Allows a Stereodirected Access to 2- and 3-Aminosugars

Stefania Mirabella,^{a,*} Francesca Cardona^a and Andrea Goti^a

^a Department of Chemistry, University of Florence, Via della Lastruccia 13, Sesto Fiorentino, FI, Italy
Email: stefania.mirabella@unifi.it

The aminohydroxylation reaction is an extremely useful method for the oxidation of alkenes to form 1,2-aminoalcohols. This moiety is present in important synthetic pharmaceutical targets like aminosugars, key constituents of a wide variety of natural substances.¹ In 2001, Donohoe and co-workers introduced an innovative method to obtain vicinal aminoalcohols through an intramolecular aminohydroxylation of olefins, called "tethered aminohydroxylation".² This method is an adaptation of the Os-mediated Sharpless aminohydroxylation, but it guarantees a full regiocontrol by tethering the nitrogen source to the allylic alcohol. We describe herein the results of the application of this method on d-glucal and d-galactal and on their derivatives, in order to obtain 2-amino and 3-aminosugars with different configurations at the stereogenic carbon atoms depending on the starting glycals. Glucals and galactals showed complementary reactivity in dependence of the stage at which the reaction was performed, i.e., directly or after double-bond shift consequent to a Ferrier rearrangement. This method allows access to both classes of 2-amino (compound **1-2**) and 3-amino (compound **3**) sugar derivatives (Figure 1). The oxazolidinone moiety of the compound **3** was easily removed with a basic treatment to obtain the free 3-aminosugar with excellent yield.




Figure 1.

References

- [1] Rai, R.; McAlexander, I.; Chang, C. T. *Org. Prep. Proced. Int.* **2005**, 37, 337.
 [2] (a) Donohoe, T. J.; Johnson, P. D.; Helliwell, M.; Keenan, M. *Chem. Commun.* **2001**, 2078; (b) Donohoe, T. J.; Callens, C. K. A.; Lacy, A. R.; Winter, C. *Eur. J. Org. Chem.* **2012**, 655.
 [3] Mirabella, S.; Cardona, F.; Goti, A. *Org. Lett.* **2015**, 17, 728-731.

Copper(II)-Catalyzed Cyclizations of Alkynes as an Efficient Tool for the Synthesis of Haloalkylidene-Substituted Heterocycles

Tea Borelli,^{*a} Gianluigi Broggin, ^a Daria Diamante^a

^a Dipartimento di Scienza e Alta Tecnologia, Università degli Studi dell'Insubria, via Valleggio 9, 22100 Como, Italy.
Email: tea_borelli@yahoo.it

Nitrogen- and oxygen-containing heterocycles are important motifs in many biologically active compounds. Intramolecular transition metal-catalyzed aminations and alkoxylation of carbon-carbon multiple bonds represent one of the most effective approaches for their synthesis.¹ Among the metals, copper is challenging because of the low cost and the tolerance toward many reactive functional groups, not requiring rigorously anaerobic and anhydrous conditions.² As a part of our ongoing interest toward intramolecular transition metal-catalyzed reactions involving C-H and heteroatom-H functionalization,³ here we describe domino copper(II)-catalyzed alkoxyhalogenation and aminohalogenation processes of alkynylureas, secondary amides and carbamates to afford a range of haloalkylidene-substituted heterocycles in good yields (Figure 1).




Figure 1.

Although the selected substrates contain a CONH group which can provide either C-N or C-O bonds, totally selective pathways were observed, depending solely on the type of substrates. The target were achieved working with stoichiometric or catalytic copper(II)-salts, in the latter case combined with N-halosuccinimide.

References

- [1] (a) Fujino, D.; Hayashi, S.; Yorimitsu, H.; Oshima, K. *Chem. Commun.* **2009**, 5754; (b) Muñiz, K.; Iglesias, A.; Fang, Y. *Chem. Commun.* **2009**, 5591.
 [2] Ueda, S.; Nagasawa, H. *Angew. Chem., Int. Ed.* **2008**, 47, 6411.
 [3] (a) Broggin, G.; Beccalli, E. M.; Borelli, T.; Brusa, F.; Gazzola, S.; Mazza, A. *Eur. J. Org. Chem.*, **2015**, 4261; (b) Broggin, G.; Poli, G.; Beccalli, E. M.; Brusa, F.; Gazzola, S.; Oble, J. *Adv. Synth. Catal.* **2015**, 357, 677.

Synthesis and Characterization of New Ruthenium(II) Arene Acylpyrazolone Complexes

Jessica Palmucci,^{a*} Fabio Marchetti,^a Francesca Condello,^b Agnese Petrini,^b Riccardo Pettinari,^b Nello Mosca,
Claudio Pettinari^b

^a School of Science and Technology, Via S. Agostino 1, 62032 Camerino (MC), Italy.

Email: jessica.palmucci@unicam.it

^b School of Pharmacy, Via S. Agostino 1, 62032 Camerino (MC), Italy

Arene ruthenium(II) complexes have been widely studied in the recent years, and their chemistry has been extensively studied for arenes such as benzene and para-cymene. They have found applications in catalysis, supramolecular assemblies, molecular devices.¹ Moreover, they have shown antiviral, antibiotic and anticancer activities.²

Recently, mononuclear Ru-acylpyrazolone complexes have been shown to exhibit interesting *in vitro* activity in some cancer cells.³

On the basis of our previous work,⁴ we extend our investigation to the interaction of several acylpyrazolones ligands with (arene)ruthenium(II)dichloride acceptors and PTA (1,3,5-triaza-7-phosphaadamantane)(Figure 1). Their structures have been confirmed by analytical and spectral data. X-ray diffraction studies and their biological activity are under investigation.




Figure 1.

References

- [1] Trost, B. M.; Frederiksen, M. U.; Rudd, M. T. *Angew. Chem. Int. Ed.* **2005**, *44*, 6630.
- [2] Yan, Y. K.; Melchart, M.; Habtemariam, A.; Sadler, P. J. *Chem. Commun.* **2005**, 4764.
- [3] Pettinari, R.; Marchetti, F.; Pettinari, C.; Petrini, A.; Skelton, B.W.; White, A.H.; Bonfili, L.; Cuccioloni, M.; Mozzicafreddo, M.; Ceardini, V.; Angeletti, M.; Nabissi, M.; Eleuteri, A.M. *J. Med. Chem.* **57**, **2014**, 4532.
- [4] Pettinari, R.; Pettinari, C.; Marchetti, F.; Petrini, A.; Skelton, B.W.; White, A.H.; Bonfili, L.; Cuccioloni, M.; Eleuteri, A.M. *J. Organomet. Chem.* **791**, **2015**, 1.

The Influence of Structure of Dialkylindium Alkoxides and Aryloxides on its Reactivity Towards N-Heterocyclic Carbenes (NHC).


Martyna Cybularczyk,^{a,b*} Paweł Horeglad^b

^a University of Warsaw, Faculty of Chemistry, L. Pasteura 1, 02-093 Warsaw, Poland.

Email: mcybularczyk@chem.uw.edu.pl

^b Centre of New Technologies, University of Warsaw, S. Banacha 2c, 02-097 Warsaw, Poland.

Despite the fact that transition-metal complexes with N-heterocyclic carbenes (NHC) have been widely used as catalysts in many different areas of chemistry, there are much fewer reports on main-group-metal complexes with NHC^{1,2}. As a part of our research, we have demonstrated that monomeric complexes Me₂Ga(NHC)(OR) - a products of reactions of [Me₂GaOR]₂ with NHC, are formed by breaking the Ga₂O₂ bridges followed by formation of strong Ga-C_{NHC} bond. Such compounds are able to polymerize racemic lactide (*rac*-LA) in controlled and stereoselective manner leading to isotactic poly(lactide) (PLA)^{3,4}. Therefore, we have been interested in synthesis, structure and catalytic activity of indium analogues. We have shown that there is a significant difference in reactivity of [Me₂InOR]₂ towards NHC, which can result in either disproportionation reaction leading to Me₃In(NHC) and Me₆In₄(OR)₆, or the formation of Me₂In(NHC)(OR) complexes (Scheme 1). On the basis of the X-ray structures and NMR spectra I am going to focus on the influence of alkoxide and aryloxy groups on the bridging In₂O₂ bonds and relate it to the reactivity of [Me₂InOR]₂ towards NHC.



Scheme 1.

References

- [1] See for example: Glorius, F.; Ed. *N-heterocyclic Carbenes in Transition-Metal Catalysis*; Springer: Berlin, Heidelberg, **2007**.
- [2] Willans, C., E. *Organometallic Chemistry*; The Royal Society of Chemistry: London, **2010**, *36*, 1.
- [3] Horeglad, P.; Szczepaniak, G.; Dranka, M.; Zachara, J. *Chem. Commun.*, **2012**, *48*, 1171.
- [4] Horeglad, P.; Cybularczyk, M.; Trzaskowski, B.; Żukowska, G., Z., Dranka, M.; Zachara, J. *Organometallic*, in press.

Sensitising NMR: Developing Magnetisation Transfer Catalysis

Amy J. Ruddlesden,^a Ryan E. Mewis^b and Simon B. Duckett^{a*}

^aCentre for Hyperpolarisation in Magnetic Resonance (CHyM), York Science Park, University of York, Heslington, York, YO10 5NY, UK,

*simon.duckett@york.ac.uk

^bSchool of Science and the Environment, Division of Chemistry and Environmental Science, Manchester Metropolitan University, John Dalton Building, Chester St., Manchester, M1 5GD, UK

Novel bidentate iridium carbene complexes have been synthesised, characterised and shown to be effective catalyst precursors for the hyperpolarisation of heterocycles. Hyperpolarisation is the method that changes insensitive NMR into a highly sensitive technique.¹ The Signal Amplification By Reversible Exchange (SABRE)² process involves the catalytic transfer of magnetisation from *para*-hydrogen (*p*-H₂) to a substrate at low magnetic field. The best performing catalysts to date are charged species that consequently fail to deliver good activity in non-polar solvents.

This research set out to address this, by synthesising novel neutral iridium complexes, **1** and **2**, and analysing their subsequent reactions with *p*-H₂ and substrate (**Errore. L'origine riferimento non è stata trovata.**). Both species efficiently enhance the NMR signals of tested substrates, in non-polar and polar solvents. Intermediate alkene dihydride complexes were observed at low temperature and the mechanism of exchange has been extensively studied.³ The active catalytic species, **3**, was found to be solvent dependent, with the Ir-O bond being stable in DCM but cleaved in methanol. For **4** this was not the case.³ Both active complexes undergo substrate and H₂ exchange thus enabling them to act as SABRE catalysts. This study provides insight into the design of effective homogeneous catalysts for magnetisation transfer.




Figure 1.

References

- [1] Lloyd, L. S.; Asghar, A.; Burns, M. J.; Charlton, A.; Coombes, S.; Cowley, M. J.; Dear, G. J.; Duckett, S. B.; Genov, G. R.; Green, G. G. R.; Highton, L. A. R.; Hooper, A. J. J.; Khan, M.; Khazal, I. G.; Lewis, R. J.; Mewis, R. E.; Roberts, A. D.; Ruddlesden, A. J., *Catal. Sci. Technol.*, **2014**, *4*, 3544.
- [2] Adams, R. W.; Aguilar, J. A.; Atkinson, K. D.; Cowley, M. J.; Elliott, P. I. P.; Duckett, S. B.; Green, G. G. R.; Khazal, I. G.; Lopez-Serrano, J.; Williamson, D. C., *Science*, **2009**, *323*, 1708.
- [3] Ruddlesden, A. J.; Mewis, R. E.; Green, G. G. R.; Whitwood, A. C.; Duckett, S. B. *Organometallics*, **2015**, *34*, 2997.

Easy-to-Synthesise, Robust, Organo-Osmium Asymmetric Transfer Hydrogenation Catalysts

J. P. C. Coverdale,^a C. Sanchez-Cano,^a I. Romero-Canelon,^a M. Wills,^{*a} P. J. Sadler^{*a}

^a Department of Chemistry, University of Warwick, Coventry, CV4 7AL (UK)

Email: j.p.c.coverdale@warwick.ac.uk

Asymmetric transfer hydrogenation (ATH) is an important process in organic synthesis for which the Noyori-type Ru^{II} catalysts [(arene)Ru(Tsdiimine)] are well established and widely used.¹ We demonstrate for the first time the catalytic activity of the osmium analogues.² X-ray crystal structures of the 16 electron Os^{II} catalysts are almost identical to those of Ru^{II}.³ The Os^{II} catalysts are readily synthesised and exhibit excellent enantioselectivity in ATH reactions (Figure 1).¹ The complexes exhibit moderate antiproliferative activity in cancer cells, which can be further enhanced by the co-administration of sodium formate.




Figure 1.

References

- [1] Reviews on asymmetric transfer hydrogenation. a) R. Noyori, S. Hashiguchi, *Acc. Chem. Res.* **1997**, *30*, 97-102. b) S. E. Clapham, A. Hadzovic, R. H. Morris, *Coord. Chem. Rev.* **2004**, *248*, 2201-2237. c) S. Gladiali, E. Alberico, *Chem. Soc. Rev.* **2006**, *35*, 226-236. d) T. Ikariya, K. Murata, R. Noyori, *Org. Biomol. Chem.* **2006**, *4*, 393-406. e) T. Ikariya, A. J. Blacker, *Acc. Chem. Res.* **2007**, *40*, 1300-1308. f) C. Wang, X. Wu, J. Xiao, *Chem. Asian J.* **2008**, *3*, 1750-1770. g) A. Robertson, T. Matsumoto, S. Ogo, *Dalton Trans.* **2011**, *40*, 10304-10410. h) J. Vaclavik, P. Sot, B. Vilhanova, J. Pechacek, M. Kuzma, P. Kacer, *Molecules* **2013**, *18*, 6804-6828.
- [2] J. P. C. Coverdale, C. Sanchez-Cano, G. J. Clarkson, R. Soni, M. Wills, P. J. Sadler, *Chem. Eur. J.* **2015**, *21*, 8043.
- [3] H. J. Haack, S. Hashiguchi, A. Fujii, T. Ikariya, R. Noyori, *Angew. Chem., Int. Ed.* **1997**, *36*, 285.

Poly(4-Vinylpyridine) Wrapped Carbon Nanotubes: an Efficient Supporting System for the Heterogenization of Methyltrioxorhenium

Issam Abdalghani,^{a*} Marcello Crucianelli,^a Bruno Bizzari,^b Giorgia Botta,^b
Davide Piccinino,^b Raffaele Saladino^b


^aDepartment of Physical and Chemical Sciences (DSFC), via Vetoio-Coppito 1, I-67100 L'Aquila,
Email: esam.ghani1988@hotmail.com

^bDepartment of Biological and Ecological Sciences (DEB), via san Camillo de Lellis, I-01100 Viterbo

Carbon nanotubes (CNTs) are well explored quasi one-dimensional nanostructured materials that are characterized by unique electronic, mechanical and structural properties. Thus far, CNTs have been proven to have great potential, among other, in material sciences and medicinal chemistry. In addition, CNTs are chemically stable in many media, present a high surface area and can undergo chemical functionalizations thus rendering them potentially very useful for catalytic ends.¹

Taking into account these aspects and following our ongoing interest in developing new reliable, inexpensive and easily preparable supports for the heterogenization of methyltrioxorhenium (MTO),² we decide to study, as an example, the behavior of new supporting systems obtained by simple grinding of pristine CNTs and polymers as poly(4-vinylpyridine)³ (PVP) and poly(4-vinylpyridine) *N*-oxide (PVPN) (Scheme, route B). Different strategies based on direct loading of MTO or of preformed MTO/PVP or MTO/PVPN adducts on pristine CNTs, have been also explored (Scheme, routes A and C).

Herein, the preliminary results we have obtained in the use of these new heterogeneous catalysts for the selective oxidation of organic substrates, in the presence of H₂O₂ as main oxidant, will be showed.



Scheme 1.

References


- [1] Tasis, D.; Tagmatarchis, N.; Bianco, A.; Prato, M. *Chem. Rev.* **2006**, *106*, 1105.
- [2] Crucianelli, M.; Saladino, R.; De Angelis, F. *ChemSusChem* **2010**, *3*, 524.
- [3] Hua, J.; Wang, Z.; Zhao, J.; Xu, L.; Zhang, J. I.; Li, R.; Sun, X. *J. Macromol. Sci. B* **2011**, *50*, 679.

Encapsulated rhodium catalysts for asymmetric hydroformylation

Lukas J. Jongkind,^{a*} Joost N.H. Reek^a

^aHomogeneous, Supramolecular and Bio-Inspired Catalysis, van 't Hoff Institute for Molecular Sciences, Universiteit van Amsterdam
Email: l.j.jongkind@uva.nl

Hydroformylation is one of the largest homogeneously catalyzed processes.¹ Because many side products can be formed under catalytic conditions, controlling selectivity is of key importance. The hydroformylation of unfunctionalized alkenes traditionally leads to the selective formation of the linear product. Previous research in our group led to the discovery of a supramolecular assembly capable of 1-octene hydroformylation with the branched product as the major product² (**1**), even at elevated temperatures.^{3,4} The selectively formed branched product in these reactions contains a stereogenic carbon atom, allowing for enantioselective hydroformylation, generating high value chiral aldehydes from cheap starting material. An efficient and widely applicable method for the asymmetric hydroformylation of unfunctionalized alkenes has not been found.



The goal of this research is to create a chiral supramolecular assembly for the asymmetric hydroformylation of unfunctionalized alkenes. In this work we explored chiral analogues of our previous successful supramolecular assembly by applying chiral porphyrin building blocks. Characterization and catalytic studies of the chiral supramolecular assemblies will be discussed

References

- [1] Franke, R.; Selent, D.; Börner, A. *Chem. Rev.* **2012**, *112*, 5675-5732
- [2] Slagt, V. F.; Reek, J. N. H.; Kamer, P. C. J.; van Leeuwen, P. W. N. M. *Angew. Chem.* **2001**, *113*, 4401-4404.
- [3] Besset, T.; Norman, D.W.; Reek, J.N.H. *Adv. Synth. Catal.* **2013**, *355*, 348-352.
- [4] Bocokic, V.; Kalkan, A.; Lutz, M.; Spek, A.L.; Gryko, D.T.; Reek, J.N.H. *Nat. Commun.* **2013**, *4*, 2670.

Dioxidomolybdenum(VI) Complexes of α -Amino Acids

Lorenzo Biancalana,^{a,*} Marco Bortoluzzi,^b Claudia Forte,^c Fabio Marchetti,^a Guido Pampaloni^a

^a Dipartimento di Chimica e Chimica Industriale, Università di Pisa, Via G. Moruzzi 13, 56124 Pisa (PI), Italy.

Email: lorenzo.biancalana@for.unipi.it

^b Dipartimento di Scienze Molecolari e Nanosistemi, Università Ca' Foscari Venezia, Dorsoduro 2137, 30123 Venezia (VE), Italy.


^c Istituto di Chimica dei Composti Organometallici, Consiglio Nazionale delle Ricerche, Via G. Moruzzi 1, 56124 Pisa (PI), Italy.

The Molybdenum(VI)/ α -amino acid system has aroused interest for the implications associated with the biochemical roles of its components as well as for the possibility to obtain potential catalysts for environmentally friendly stereoselective reactions.¹

Studies on the interaction between molybdates and α -amino acids in water have been mostly devoted to α -amino acids with a polar or ionisable side-chain and only in few cases a well-defined product has been isolated in the solid state.²

On the other hand, the interaction between Mo(VI) and α -amino acids with a non-polar side chain has been much less investigated in solution, although solid products have been isolated from highly acidic media (pH = 0-2).^{3,4} The exploration of this piece of chemistry holds an important drawback: in general, the products are insoluble materials whose crystallographic characterization still remains a hard, not achieved task. As a matter of fact, only limited and often controversial spectroscopic data have been reported so far, therefore the Mo-oxido scaffold and the amino acid coordination mode have not been clearly elucidated.

In the light of this preface, we investigated the reactions of Na₂MoO₄·2H₂O and (NH₄)₆Mo₇O₂₄·4H₂O with a selection of α -amino acids in acidic aqueous solution and carried out a detailed solid state characterization of the isolated Mo(VI) complexes.⁴ The combination of IR and NMR data and DFT calculations has allowed the unprecedented structural identification of the dinuclear core and of the $\kappa(O,O')$ bidentate bridging coordination mode of the α -ammonium acidate ligand. The formation of coordination polymers presumably occurs by condensation reactions.



References

- [1] Severin, K.; Bergs, R.; Beck, W. *Angew. Chem. Int. Ed.* **1998**, *37*, 1634.
- [2] a) Butcher, R.J.; Powell, H.K.J.; Wilkins, C.J.; Yong, S.H. *J. Chem. Soc., Dalton Trans.* **1976**, 356. b) Kay, A.; Mitchell, P.C. *J. Chem. Soc. A* **1970**, 2421.
- [3] Eguchi, H.; Takeuchi, T.; Ouchi, A.; Furuhashi, A. *Bull. Chem. Soc. Jpn.* **1969**, *42*, 3585.
- [4] Biancalana, L.; Bortoluzzi, M.; Forte, C.; Marchetti, F.; Pampaloni, G. *RCS Adv.* **2015**, *5*, 9010 and references therein.

Photophysical Properties of Zn-Porphyrin Complexes Decorated with an Easily Accessible Anchoring Ligand

Paolo Boggio,^{a,b*} Annamaria Deagostino,^a Teodor Silviu Balaban^b

^a Dep. of Chemistry, University of Torino, Via P. Giuria 7, 10125 Torino (TO), Italy.

Email: paolo.boggio@unito.it

^b Aix Marseille Université, CNRS UMR 7313, Institut des Sciences Moléculaires de Marseille (iSm2) Chirosciences, Service 442, Avenue Escadrille Normandie-Niemen, F-13397 Marseille CEDEX 20, France.

Since their discovery, porphyrins have played an important role in the organic and organo-metallic chemistry. They have been studied as dyes in energy production devices, like dye sensitized solar cells (DSSC),¹ or as organic ligands for metallo-complexes in photosynthetic systems, inspired by natural light harvesting complexes.² Carboxylic groups are usually used for anchoring dyes on to semiconductor surface.³ Here we present the results of the photophysical studies for two different Zn-porphyrin complexes synthesized starting from a new, versatile porphyrin ligand. (Fig 1)




Figure 1. UV-Vis spectra for the two Zn-complexes [2] and [3].

Zn-porphyrinate complex [2] shows self-assembly in non-coordinating solvents like *n*-heptane and can be easily functionalized exploiting the porphyrin free *meso* position and the carboxylic group at the end of the linear chain. These aspects make it a potential and versatile building block for supramolecular architectures.

Zn-porphyrinate complex [3] presents a wider wavelength range for light absorption than Zn-complex [2]. Furthermore its generation occurs in very soft experimental conditions and doesn't require metallic catalysts [like Ag (I)] usually employed to promote the *meso-meso* coupling reaction between metallo-porphyrins.⁴

References

- [1] Shahroosvand, H.; Zakavi, A.; Sousaraei, A.; Eskandari, M. *Phys. Chem. Chem. Phys.* **2015**, *17*, 6347.
- [2] Kelle, R. F.; Tauber, M. J.; Wasielewski, M. R. *Angew. Chem. Int. Ed.* **2006**, *45*, 7979.
- [3] Mathew, S.; Yella, A.; Gao, P.; Humphry-Baker, R.; Curchod, B. F. E.; Ashari-Astani, N.; Tavarnerelli, I.; Ro Rothlisberger, U.; Nazeeruddin, M. K.; Gratzel, M. *Nat. Chem.* **2014**, *6*, 242
- [4] Maeda, C.; Kamada, T.; Aratani, N.; Osuka, A. *Coord. Chem. Rev.* **2007**, *251*, 2743.

Cage Controlled Catalysis through Co-Encapsulation of Catalyst and Substrate

Sandra S. Nurttila^a and Joost N.H. Reek^{a*}

^a Homogeneous, Supramolecular and Bio-inspired Catalysis, van 't Hoff Institute for Molecular Sciences, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands, Email: j.n.h.reek@uva.nl, s.s.nurttila@uva.nl

Nature arrives at high selectivities and activities in enzymatic transformations by confining the active site in a bulky protein structure. Due to the complexity of proteins, simpler, synthetically more accessible cage analogues have been studied to ultimately achieve higher activity and selectivity in known catalytic transformations. Our group has previously concluded that the co-encapsulation of substrate and catalyst within synthetic cages or the template-ligand approach are two of the potential strategies for nature-inspired catalysis in confined spaces (see figure below).¹ In this study we explore the co-encapsulation of various guests and catalysts within synthetic cages, to evaluate the potential of this approach using this specific type of cage compounds. We evaluate the concept of co-encapsulation within synthetic M₄L₆ capsules designed by the group of Nitschke.²⁻⁴ Larger porphyrin-edged capsules designed by the same group are currently also being explored.⁵




Figure 1.

This poster will present the recent results of the co-encapsulation of different transition metal complexes, ligands and substrates within cationic tetrahedral capsules constructed of bis-bidentate pyridyl-imine based ligands coordinated to iron(II) vertices. Spectroscopic data confirming the encapsulation of various guests within the synthetic cages will be demonstrated. Preliminary results on the activity of encapsulated species in catalytic transformations will be discussed as well.

References

- [1] Leenders, S.H.A.M.; Gramage-Doria, R.; de Bruin, B.; Reek, J.N.H. *Chem. Soc. Rev.* **2015**, *44*, 433.
- [2] Ronson, T.K.; Zarra, S.; Black, S.P.; Nitschke, J.R. *Chem. Commun.* **2013**, *49*, 2476.
- [3] Bolliger, J.L.; Belenguer, A.M.; Nitschke, J.R. *Angew. Chem. Int. Ed.* **2013**, *52*, 7958.
- [4] Mal, P.; Breiner, B.; Rissanen, K.; Nitschke, J.R. *Science* **2009**, *324*, 1697.
- [5] Wood, D.M.; Meng, W.; Ronson, T.K.; Stefankiewicz, A.R.; Sanders, J.K.M.; Nitschke, J.R. *Angew. Chem. Int. Ed.* **2015**, *54*, 3988.




Synthesis and Functionalization via Direct Lithiation of α -Aryl-substituted Tetrahydropyrans

Luciana Cicco,^{a*} Francesca Claudia Sassone,^a Antonio Salomone,^a Filippo Maria Perna,^a Carsten Adam Donau,^c Konstantin Karaghiosoff,^c Andrea Temperini,^b Vito Capriati^a

^a Dipartimento di Farmacia-Scienze del Farmaco, Università di Bari "Aldo Moro", Consorzio C.I.N.M.P.I.S., Via E. Orabona 4, I-70125 Bari, Italy.

^b Dipartimento di Scienze Farmaceutiche, Università di Perugia, Via A. Fabretti 48, I-06123 Perugia, Italy. ^c Department of Chemistry, Ludwig Maximilians University, Munich, Butenandtstrasse 5-13, Munich 81377, Germany. e-mail: luciana.cicco@uniba.it

Many biologically active and widely distributed natural products contain a tetrahydropyran moiety. The tetrahydropyran core is, indeed, a useful "building block" not only in organic chemistry, but also in biological and medicinal chemistry.¹ Building on our recent successful α -lithiation-electrofilic interception protocols applied to phenyloxetane² and phenyltetrahydrofuran³ derivatives, we have now extended the result of such an investigation to the preparation of 2,2-disubstituted tetrahydrofurans **2** starting from the parent α -arylated precursors **1** (Scheme 1). The synthesis of racemic **3**, starting α -aryl-substituted tetrahydropyrans has been accomplished exploiting a one-pot strategy, under microwave conditions, which involves a preliminary Heck cross-coupling reaction followed by an hydrogenation procedure (Scheme 2). On the other hand, the preparation of the corresponding optically active substrates was based on a selenium-mediated cyclization starting from δ -phenylseleno ketones (Scheme 3).² In this communication, the influence of the solvent and the temperature, and the nature of the organolithium employed, on the effectiveness of such a direct α -functionalization of these oxygenated heterocycles will be discussed as well as the chemical and configurational stability of the putative anion **1-Li** (Scheme 2).



Acknowledgements: This work was financially supported by the Interuniversities Consortium C.I.N.M.P.I.S., and by the University of Bari.

References

- [1] N. M. Nasir, K. Ermanis, P. A. Clarke, *Org. Biomol. Chem.* **2014**, *12*, 3323.
- [2] D. I. Coppi, A. Salomone, F. M. Perna, V. Capriati, *Chem. Commun.* **2011**, *47*, 9918.
- [3] R. Mansueto V. Mallardo, F. M. Perna, F. Sassone, A. Salomone, V. Capriati, *Chem. Commun.* **2013**, *49*, 10160.
- [4] A. Temperini, A. Barattucci, P. M. Bonaccorsi, O. Rosati, L. Minuti, *unpublished results*.


Aminoethyl Glycine Peptide Nucleic Acids (*aeg*-PNA): the Study of their Interaction with DNA by Means of Stopped Flow Technique

Pramod R. Thakare,^{a*} Silvia Cauteruccio,^a Claudio Carrara,^a Emanuela Licandro,^a Nadia Barbero,^b Guido Viscardi^b

^a Department of Chemistry, University of Milan, Via C. Golgi 19, 20133-Milan, Italy. e-mail: pramod.thakare@unimi.it

^b Department of Chemistry, University of Turin, Via Giuria 7, 10125-Turin, Italy.

Stopped-flow technique is useful for studying fast reactions that have half-lives as short as a few milliseconds.¹ This technique has never been used to study the interaction between Peptide Nucleic Acids (PNA) and DNA strands in the formation of hybrids. PNAs are mimics of DNA, in which the entire sugar-phosphate backbone has been replaced by a peptide amide bond backbone composed of *N*-(2-aminoethyl) glycine subunits. PNAs show unique properties that set them apart from other traditional DNA analogues.² Normally the affinity of PNA for DNA is detected by using melting temperature³ but sometimes this technique is not appropriate, especially if PNA contain some thermal sensitive functional groups. So, we have started an innovative study of PNA-DNA interaction by using stopped-flow technique and for this we have synthesised the PNA sequences reported in Scheme 1.



Scheme 1. Synthesis of *aeg*PNA decamers.

References

- [1] Harvey R. A.; Borcherdt W. O. *Anal. Chem.* **1972**, *44*, 1926-1928.
- [2] Hyrup B.; Nielsen P. E. *Bioorg. Med. Chem. Lett.* **1996**, *4*, 5-13.
- [3] Schwarz F. P.; Robinson S.; Butler J. M., *Nucleic Acids Research*, **1999**, *27*, 4792-4800.

Epoxidation Reactions in a Self-Assembled Molecular Cage

Petrus F. Kuijpers,^{a*} Joost N. H. Reek,^a Bas de Bruin^a

^a Homogeneous Supramolecular and Bio-Inspired catalysis, Van 't Hoff institute for molecular Sciences, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands.

*P.F.Kuijpers@uva.nl

Both in industry and academia obtaining high activity and selectivity is required for applied catalysts. Although many highly active and selective catalysts have been reported to date, many reactions still suffer from low activity, selectivity or even both. As in nature's catalysts, enzymes, applying a second coordination sphere could supply the solution for these challenging reactions.¹ In our research we investigated the use of a self-assembled molecular cage to improve the catalytic performance of various catalysts.

The cubic M_8L_6 cage depicted in figure 1 has been shown to be suitable for the encapsulation of pyridine functionalized porphyrins.² The Zn-porphyrin planes of this cubic cage can be used to selectively coordinate catalysts containing a pyridine moiety. Recently we reported the possibility to perform size-selective catalysis within the confinement of this molecular cage with encapsulated cobalt porphyrins.³




Figure 1.

The main degradation pathways for manganese porphyrin catalysts in oxidation reactions are bimetallic. Therefore we envisioned that encapsulation of a manganese catalyst would improve the lifetime and therefore the activity of the catalyst. In this poster we will disclose the results of our research towards improving the catalytic activity of a manganese porphyrin by encapsulation in a molecular cage. Secondly a cage induced substrate selectivity will be discussed.

References

- [1] Leenders, S.H.A.M.; Gramage-Doria, R.; de Bruin, B.; Reek, J.N.H. *Chem. Soc. Rev.* **2015**, *44*, 433.
- [2] Otte, M.; de Bruin B. et al, *Chem. Eur. J.* **2013**, *19*, 10170.
- [3] Otte, M.; de Bruin B. et al, *Chem. Eur. J.* **2014**, *20*, 4880.

Microwave-Assisted Transesterification and Hydroformylation of Natural Oils: a Sustainable Approach for Transformation of *Calophyllum inophyllum* into Value-Added Products

Liliana Damas^a, Gonçalo N. Costa^{a,b}, Juvêncio C. Ruas^a, Rui M. B. Carrilho^a, Gilberto Aquino^c, Mário J.F. Calvete^a, Marta Pineiro^{a*} and Mariette M. Pereira^{a*}

^a CCC, Department of Chemistry, University of Coimbra, Rua Larga, 3004-535 Coimbra, Portugal. ^b Luzitin, SA, Edifício Bluepharma, São Martinho do Bispo, 3030-016 Coimbra, Portugal. ^c Faculty of Pharmacy, Exact Sciences and Technology Unit, State University of Goiás, BR-75132400 Anápolis, Goiás, Brazil.

The use of renewable resources in the manufacturing of value-added products is one of the great challenges for the chemical researchers of this decade.¹ The direct use of vegetable oils and animal fats as combustible fuels is not a suitable alternative to fossil fuel oils due to their high viscosity and low volatility. However the transesterification reaction of natural oils allows the reduction of these drawbacks and therefore it has been commonly used for the production of biodiesel, including the use of catalysts based on renewable sources.² Furthermore, the transition metal catalyzed hydroformylation reaction constitute an efficient process to promote the reduction of high olefin content in an one-step transformation of natural unsaturated fatty esters into the corresponding aldehydes, or derivatives, thereby increasing their economic value. Recently, the use of microwave (MW) irradiation to promote transesterification reactions of natural oils has attracted the attention of several researchers due to the demands for the development of more energetically friendly processes.^{2,3} In this work we describe an eco-friendly approach for the transesterification of *Calophyllum inophyllum* seeds oil, based in the use of heterogeneous catalysts obtained from renewable sources under microwave irradiation, followed by hydroformylation of the resultant long chain methyl esters.




Figure 1. (A) Microwave assisted reactions of natural oils; (B) CEM Discover® SP microwave synthesizer connected to a syngas addition system.

References


- [1] Karmarkar, A.; Karmarkar, S.; Mukherjee, S. *Bioresour. Technol.*, **2010**, *101*, 7201.
- [2] Issariyakul, T.; Dalai, A.K. *Renew. Sust. Ener. Rev.*, **2014**, *31*, 446.
- [3] Damas, L.; Costa, Gonçalo N.; Ruas, Juvêncio C.; Carrilho, R. M. B.; Abreu, Artur R.; Aquino, G.; Calvete, M. J. F.; Pineiro, M.; Pereira, Mariette M. *Current Microwave Chemistry*, **2015**, *2*, 53.

Role of Electron Transfer Processes in the *N*-demethylation of *N,N* dimethylanilines and *S*-Oxidation of Aromatic Sulfides Promoted by the Nonheme Iron Complex $[\text{Fe}^{\text{II}}(\text{N4Py})](\text{OTf})_2$

Alessia Barbieri,^{a*} Osvaldo Lanzalunga^a

^a Dipartimento di Chimica and Istituto CNR di Metodologie Chimiche-IMC, Sezione Meccanismi di Reazione, Università di Roma "La Sapienza", P.le A. Moro 5, 00185 Rome, Italy

The role of electron transfer processes has been investigated in two of the most important reactions promoted by nonheme iron(II) complexes, namely the *N*-demethylation of *N,N*-dimethylanilines and *S*-Oxidation of Aromatic Sulfides. In the oxidative *N*-demethylation of *N,N*-dimethylanilines promoted by nonheme iron(IV)-oxo species, a mechanistic dichotomy between a direct hydrogen atom transfer and a sequential electron transfer/proton transfer mechanism (ET/PT) can be envisaged (Scheme 1).^{1,2}



Scheme 1.





Figure 1

We have investigated the kinetic isotope effects profiles in the *N*-demethylation of a series of 4-*X*-*N,N*-dimethylanilines promoted by $[\text{Fe}^{\text{IV}}(\text{O})(\text{N4Py})]^{2+}$ (Figure 1). The results obtained are in agreement with the ET/PT mechanism in which the deprotonation of the anilinium radical cation represents the rate determining step. The bell shaped KDIE_{intra} vs. pKa anilinium radical cation profile observed, allowed us to estimate the pKa value of $[\text{Fe}^{\text{III}}(\text{OH})(\text{N4Py})]^{2+}$. In the oxidation of aromatic sulfides to sulfoxides catalyzed by nonheme iron(IV) oxo complex, two mechanisms are possible: direct oxygen atom transfer (Scheme 2, path a) and a sequential electron transfer/oxygen transfer mechanism (Scheme 2, path b-c).^[3] Using aryl diphenylmethyl sulfides whose radical cations are characterized by high fragmentation rate constants, we found that the formation of the sulfoxide is accompanied by products deriving from the α -C-S and α -C-H bond cleavage in the radical cation (Scheme 2, path d,e) thus supporting the occurrence of the ET-oxygen rebound mechanism.



Scheme 2.

References

- [1] Nam W., Lee Y.-M., Fukuzumi S. *Acc. Chem Res.* **2014**, *47*, 1146–1154.
- [2] Nam W. *Acc. Chem Res.* **2007**, *40*, 522–531.
- [3] Park J., Morimoto Y., Lee Y.-M., Nam W. *J. Am. Chem. Soc.* **2011**, *133*, 5236–5239.

Poster 33

Synthesis of Arylated Olefins Catalyzed by Palladium(II) Complexes

Ewelina Silarska,^{a*} Anna M. Trzeciak^a

^a University of Wrocław, Faculty of Chemistry, Joliot –Curie 14, 50- Wrocław, Poland

Email: ewelina.silarska@chem.uni.wroc.pl

Palladium-catalyzed cross-coupling reaction is a powerful tools for synthesis of arylated olefins. Arylated olefins are ubiquitous structural motifs in natural products, pharmaceutical intermediates, cosmetology and organic materials.¹ Of the many organometalloids known to undergo coupling reactions, organosilicon reagents are the most intriguing since they are inexpensive, non-toxic, readily prepared and environmentally friendly.




Figure 1. Reaction of organosilanes with olefins catalyzed by anionic palladium complexes.

In this work we present a new way leading to arylated olefins, based on a cross-coupling triphenylsilane and triphenylsilanol with different olefins catalyzed by anionic palladium(II) complexes (Fig.1).² The reaction was studied with variation of parameters such as different oxidants (e.g. copper salts, oxygen, AgF), salts, time and solvents. Studies shown a positive effect of the presence of pyridinium cations on the catalytic activity of palladium complex. Also effect of the temperatures on the selectivity of the reaction have been observed.

References

- [1] Mi X., Huang M., Gou H., Wu Y., *Tetrahedron*, **2013**, 69, 5123.
- [2] Silarska E., Trzeciak A.M., Skrzypczak A., Pernak J., *Applied Catalysis. A: General*, **2013**, 466, 216.

Poster 34

Development of simple and mild hydroamination reaction catalyzed by NHC-gold complexes

Malina Michalska,^{a*} Karol Grela,^{a*}


Biological and Chemical Research Centre, Faculty of Chemistry, University of Warsaw, Zwirki iWigury 101, 02-089 Warsaw, Poland.

Email: mmichalska@cnbc.uw.edu.pl, karol.grela@gmail.com

An hydroamination reaction of 2-alkynylanilines as substrates is potentially an important transformation. It can give rise to indole derivatives amenable to further interesting synthetic manipulations en route to natural products and some analogues.¹⁻²

However, there are only few reports describing the hydroamination which have been published to date.³⁻⁵ Various sources of catalysts, including palladium, platinum, as well as copper were used in heterocyclizations, and relatively drastic conditions were systematically employed.⁶⁻⁸ Thus, the development of an efficient and general catalytic approach for these reactions is highly desirable.

The project is devoted to the development of the simple and mild method for synthesis of indoles. Preliminary results establishing the feasibility of the approach will be presented. (Scheme 1)



Scheme 1.

References

- [1] H. L. Pearce A. Brossi, M. Suffness; In *The Alkaloids* Academic: San Diego, 1990; Vol. 37, p 145.
- [2] U. Anthoni, C. Christophersen, P. H. Nielsen In *Alkaloids: Chemical and Biological Perspectives*; Pelletier, S. W., Ed.; Wiley: New York, 1999; Vol. 13, p 163.
- [3] L. Ackermann, *Org. Lett.*, **2005**, 7, 439.
- [4] T. Shimada, I. Nakamura, Y. Yamamoto *J. Am. Chem. Soc.*, **2004**, 126, 10546.
- [5] D. Janreddy, V. Kavala, Ch.W. Kuo, T. S. Kuo, Ch. H. He *Tetrahedron*, **2013**, 69, 3323.
- [6] X. Deng, H. McAllister, N. S. Mani *J. Org. Chem.*, **2009**, 74, 5742.
- [7] T. Shimada, I. Nakamura, Y. Yamamoto *J. Am. Chem. Soc.* **2004**, 126, 10546.
- [8] R. Álvarez, C. Martínez, Y. Madich, J. G. Denis, J. M. Aurrecochea, Dr. Á. R. de Lera *Chem.-Eur. J.* **2010**, 16, 12746.

Switching Selectivity of Gold Catalysts by Encapsulation in a Self-Assembled Cage

Anne C. H. Jans,^{a*} Adrián Gómez-Suárez,^b Steven P. Nolan,^b Joost N. H. Reek^a

^a Homogeneous and Supramolecular Catalysis Group, Van 't Hoff Institute for Molecular Sciences, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands, *a.c.h.jans@uva.nl

^b Nolan Research Group, School of Chemistry, University of St Andrews, United Kingdom

Supramolecular catalysis is inspired by enzymes, which show very high activities and selectivities by using a confined space around their active center. This principle can be mimicked by encapsulation of transition metal complexes in a synthetic capsule.¹ In this research project, a self-assembled hexameric resorcin[4]arene cage is employed (Figure 1), which can encapsulate the gold carbene complex IPr-Au-OTf.²




Figure 1. A self-assembled hexameric resorcin[4]arene cage.²

In the current contribution we use encapsulation as a way to completely switch the selectivity of dinuclear gold catalysts³. Outside the capsule, the complex exists as a dimer that is able to catalyze dimerization of the substrate⁴ (Figure 2). However, the dinuclear complex is too large to be encapsulated, so inside the cage it exists as a monomeric species. As the monomeric and dimeric complexes have a completely different selectivity, the product distribution of the reaction reverses upon encapsulation.




Figure 2. The cage completely changes the selectivity of the gold catalyst.

References

- [1] Leenders, S. H. A. M.; Gramage-Doria, R.; de Bruin, B.; Reek, J. N. H. *Chem. Soc. Rev.* **2015**, *44*, 433.
- [2] Cavarzan, A.; Scarso, A.; Sgarbossa, P.; Strukul, G.; Reek, J. N. H. *J. Am. Chem. Soc.* **2011**, *133*, 2848.
- [3] Ramón, R. S.; Gaillard, S.; Poater, A.; Cavallo, L.; Slawin, A. M. Z.; Nolan, S. P. *Chem. Eur. J.* **2011**, *17*, 1238.
- [4] Hashmi, A. S. K.; Lauterbach, T.; Nösel, P.; Højer Vilhelmsen, M.; Rudolph, M.; Rominger, F. *Chem. Eur. J.* **2013**, *19*, 1058.

Mono- and Multinuclear Organometallic Rhodium and Ferrocenyl Quinolines as Potential Antiparasitic Agents

Tameryn Stringer,^{a*} Gregory S. Smith.^a

^a Department of Chemistry, University of Cape Town, Rondebosch, 7701, Cape Town, South Africa.

Email: strtam001@myuct.ac.za

One of the most frequent occurring infectious diseases worldwide is malaria. The World Health Organization (WHO) reported approximately 198 million cases of malaria in 2013.¹ One of the most successful antimalarial drugs, chloroquine (CQ), has been rendered almost useless due to the occurrence of drug resistance.² Currently, artemisinin-based combination therapy is the main treatment for malaria, but resistance in some parts of the world has recently been documented.³ Ferroquine (FQ), a ferrocene-containing derivative of CQ, shows better activity than its parent compound, CQ.⁴ Based on the success of this metal-based drug, interest has been generated towards using metal complexes for the treatment of parasitic diseases. This prompted investigation of the antiparasitic activity of mononuclear and dinuclear organometallic complexes of chloroquine (Figure 1). It is believed that increasing the number of active moieties positively influences biological activity.⁵ This study investigated the effect of incorporating polyamine scaffolds, multinuclearity as well as lipophilicity on antiparasitic activity.




Figure 1.

References

- [1] The World Health Organization, World Malaria Report 2014 http://www.who.int/malaria/media/world_malaria_report_2014/en/
- [2] Trigg P. L.; Kondrachine, A. V., In *Malaria*, I. W. Sherman, Ed., ASM Press: Washington, DC, 1998, 11.
- [3] Enserink M. *Science*, **2010**, *328*, 844.
- [4] Biot, C.; Glorian, G.; Macejewski, L. A.; Brocard, J. S.; Domarle, O.; Blampain, G.; Millet, P.; Georges, A. J.; Abessolo, H.; Dive, D. *J. Med. Chem.*, **1997**, *40*, 3715.
- [5] Stringer, T.; Taylor, D.; de Kock, C.; Guzgay, H.; Au, A.; An, S. H.; Sanchez, B.; O'Connor, R.; Patel, N.; Land, K. M.; Smith, P. J.; Hendricks, D. T.; Egan, T. J.; Smith, G. S. *Eur. J. Med. Chem.*, **2013**, *69*, 90.

Multi-Component Assemblies of tris(2-pyridylmethyl)amine Ligands, Metals and Amino Acids for Chiroptical Sensing

Nadia Alessandra Carmo dos Santos,^{a*} Elena Badetti,^a Giulia Marina Licini,^a Cristiano Zonta^a

^aDepartment of Chemical Science, Via Via Marzolo 1 - 35131 Padova (PD) Italy.
Email: nadiaalessandra.carmodossantos@studenti.unipd.it

Stereodynamic optical probes own generally a molecular fragment present in two enantiomeric forms which in presence of an analyte give rise to a preferential diastereoisomer able to furnish an optical readout.¹ Recently we reported about a new molecular probe used for the reliable determination of the enantiomeric excess of free amino acids.² Herein we report the use of new metals and the measurement of the induced circular dichroism of the resulting multicomponent assembly. The study highlight the complex equilibria present in solution for the formation of the assembly and the specie responsible of the CD signal(Figure 1).




Figure 1.

References

- [1] Leung D., Ok Kang S. and AnsynE. V. *Chem.Soc.Rev.*, **2012**,41, 448–479.
[2] Scaramuzzo, F. A.; Licini, G.; Zonta, C. *Chem. Eur. J.* **2013**, 19, 16809.

Cooperative Coordination and Ionic Interactions Assisted *E/Z* Isomerization of Arylhydrazone of Ethyl 2-Cyanoacetate in Complexation with Copper(II)

Nuno M. R. Martins,^{a*} Kamran T. Mahmudov,^{a,b} M. Fátima C. Guedes da Silva,^a

Luísa M. D. R. S. Martins,^{a,c} Armando J. L. Pombeiro^a

^a Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisbon, Portugal.
E-mail: nunommartins@tecnico.ulisboa.pt

^b Department of Chemistry, Baku State University, Z. Xalilov Str. 23, Az 1148 Baku, Azerbaijan

^c Chemical Engineering Department, ISEL, R. Conselheiro Emídio Navarro, 1959-007 Lisboa, Portugal

The *E/Z* isomerization around the double bond in organic molecules recently attracted considerable attention, mainly in view of their use as catalysts, sensors, gels, liquid crystals, molecular switches, etc.^[1-3] For instance, controllable *E/Z* isomerisation around the C=N double bond in hydrazones, what makes them good candidates for the construction of molecular switches, electronics, rotors and similar nanomachines.^[1,2] All reported *E/Z* isomerizations in hydrazones are regulated by pH, coordination-coupled proton transfer, resonance-assisted hydrogen bonding [X...H-Y ↔ Y...H-X (X, Y = N, O, S, etc.), RAHB] or its cooperation with ionic interactions, auxiliary ligand or solvation.^[1-4] Here we report a new cooperative coordination and ionic interactions assisted *E,Z* → *E* or *E,Z* → *Z* isomerization of (*E/Z*)-4-(2-(1-cyano-2-ethoxy-2-oxoethylidene)hydrazinyl)-3-hydroxybenzoic acid (**H₃L**), which occurs upon interaction with Cu^{II} in different reaction conditions (Scheme). Aqueous soluble [Cu₂(X)₂(μ-HL)₂], X = CH₃OH (**1**), (CH₃)₂NCHO (**2**) and [Cu(en)₂HL]·CH₃OH·H₂O (**3**) Cu^{II} complexes were isolated by reaction of Cu^{II} nitrate hydrate with the H₃L, in the presence (for **3**) or absence (for **1** and **2**) of ethylenediamine (en), and were characterized by elemental analysis, IR spectroscopy and X-ray single crystal diffraction.




Figure 1. *E/Z* isomerization of H₃L in the synthesis of 1–3.

References

- [1] C. Dugave, L. Demange, *Chem. Rev.* **2003**, 103, 2475.
[2] L. A. Tatum, X. Su, I. Aprahamian, *Acc. Chem. Res.* **2014**, 47, 2141.
[3] K. T. Mahmudov, M. N. Kopylovich, M. F. C. Guedes da Silva, A. J. L. Pombeiro, *ChemPlusChem* **2014**, 79, 1523
[4] K. T. Mahmudov, M. F. C. Guedes da Silva, M. Sutradhar, M. N. Kopylovich, F. E. Huseynov, N. T. Shamilov, A. A. Voronina, T. M. Buslaeva, A. J. L. Pombeiro, *Dalton Trans.* **2015**, 44, 5602.

Synthesis and Biological Evaluations of Novel Arene Ru(II) Complexes

of 4-acyl-5-pyrazolone-based Hydrazone Ligands

Francesca Condello,^{a,*} Jessica Palmucci,^b Agnese Petrini,^a Ilaria Torquati,^a Riccardo Petrelli,^a Loredana Cappellacci,^a Anders Hofer,^c Farahnaz Ranjbarian,^c A. J. L. Pombeiro,^d Giulio Lupidi,^a Riccardo Pettinari,^a Fabio Marchetti,^b Claudio Pettinari^a

^a School of Pharmacy and ^b School of Science and Technology, Via S. Agostino 1, 62032 Camerino (MC), Italy.


Email: francesca.condello@unicam.it

^c Department of Medical Biochemistry and Biophysics, Umeå University, SE-90187 Umeå, Sweden.

^d Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais 1049-001, Lisboa, Portugal.

We Hydrazone compounds, bearing –CH=N–NH–C(O)– functional groups, are a special kind of Schiff bases which have extensive biological properties. In fact their chemistry is particularly intriguing due to their capability to easily furnish valuable chemotherapeutics such as anticancer, antibacterial, antifungal and antiprotozoal agents¹ as well as enzyme inhibitors.²

As an extension of our research on acylpyrazolone chemistry³ and relative Ru-arene compounds⁴ the present work has the aim to design, synthesize and study the biological properties of new half-sandwich η^6 -arene-ruthenium(II) complexes with different 4-acyl-5-pyrazolone-based hydrazone scaffolds with general structure HL' (Figure 1). Preliminary results on DNA vs BSA protein binding and on the effect as anti-*Trypanosoma brucei* parasite agents⁵ will be reported.



HL': R¹ = N or C; R² = H or NO₂;
R³ = H or NO₂; R⁴ = Ph or Bz




Figure 1.

References

- [1] Verma, G.; Marella, A.; Shaquiquzzaman, M.; Akhtar, M.; Ali, M. R.; Alam, M. M. *J Pharm Bioall Sci* **2014**, *6*, 69.
- [2] Rupiani, S.; Buonfiglio, R.; Manerba, M.; Di Ianni, L.; Vettrai, M.; Giacomini, E.; Masetti, M.; Falchi, F.; Di Stefano, G.; Roberti, M.; Recanatini, M.; *European Journal of Medicinal Chemistry* **2015**, *101*, 63.
- [3] Marchetti, F.; Pettinari, R.; Pettinari, C. *Coordination Chemistry Reviews* **2015**, *303*, 1.
- [4] Pettinari, R.; Marchetti, F.; Pettinari, C.; Petrini, A.; Scopelliti, R.; Clavel, C. M.; Dyson, P. J. *Inorg. Chem.* **2014**, *53*, 13105.
- [5] Jacobs, R. T.; Nare, B.; Phillips, M. A. *Curr. Top. Med. Chem.* **2011**, *11*, 1255.

The bimca-ligand: Synthesis and Application of its Rhodium and Iridium Complexes

Eva Jürgens,^{a,*} Alexander Seybold,^a Doris Kunz^a

^a Institute for Inorganic Chemistry, University of Tübingen, Auf der Morgenstelle 18, 72074 Tübingen.

Email: eva.juergens@uni-tuebingen.de

Introduced by Shaw in 1976, pincer-type ligands have become an important ligand class in organometallic chemistry during the last decades.¹ Nearly 15 years ago, they were combined with NHCs either as backbone or side donor moieties. Nowadays a broad range of tridentate mono- or biscarbene ligands are increasingly used in the synthesis of highly active homogenous catalysts.² In 2007 we reported the monoanionic bis(imidazolynilidene)carbazole (bimca) ligand **2a** as anionic pincerligand with two carbene moieties.³ Since then we established a simplified synthetic access to its precursor **2** and prepared a decent-sized ligand library **2a-g**.⁴ The stable lithium and potassium carbene complexes **3,4ag** were synthesized by deprotonation with an appropriate basic precursor and could be characterized by NMR spectroscopy. Further reaction with Rh(I) or Ir(I) carbonyl precursors led to the formation of M(bimca)(CO) complexes (M = Rh, Ir) **5,6a-g**, while after the reaction with precursors such as Vaska's complex or Rh(m-Cl)(cod)]₂ also MIII(bimca) complexes were observed.^{5,7}




Figure 1.

The Rh(I) and Ir(I) carbonyl complexes **5,6a-g** show a highly nucleophilic reactivity caused by the strong σ -donor and weak π -acceptor character of the NHC moieties in addition to the monoanionic carbazolidone donor.³⁻⁶ Their nucleophilic character could be applied in different catalytic and stoichiometric reactions. Both complexes show an interesting reactivity towards allyl halides leading to the η^1 -allyl complexes.^{3,5} The Rh(I) complex **3a** shows a high catalytic activity in the *Meinwald* rearrangement of epoxides.⁶

References

- [1] (a) Pugh, D.; Danopoulos, A. *Coord. Chem. Rev.* **2007**, *251*, 610–641; (b) Albrecht, M.; van Koten, G.; *Angew. Chem. Int. Ed.* **2001**, *40*, 3750–3781.
- [2] Jürgens, E.; Kunz, D. in *Molecular Catalysts* **2014**, 183–206.
- [3] Moser, M.; Wucher, B.; Rominger, F.; Kunz, D. *Organometallics* **2007**, *26*, 1024–1030.
- [4] *Manuscript in preparation*
- [5] Seybold, A.; Wucher, B.; Alles, M.; Rominger, F.; Maichle-Mössmer, C.; Kunz, D. *J. Organomet. Chem.* **2015**, *775*, 202–208.
- [6] Jürgens, E.; Wucher, B.; Rominger, F.; Törnroos, K. W.; Kunz, D. *Chem. Commun.* **2015**, *51*, 1897–1900.
- [7] *Unpublished results*

Borate Carbene Complexes: Synthesis, Reactivity and Biological Evaluation

Marika Marinelli,^a Maura Pellei,^a Cristina Marzano,^b Valentina Gandin,^b Alessandro Dolmella,^b

H. V. Rasika Dias^c and Carlo Santini^a

^a School of Science and Technology, Chemistry Division, University of Camerino, via S. Agostino 1, 62032 Camerino, Italy.

Email: marika.marinelli@unicam.it

^b Department of Pharmaceutical and Pharmacological Sciences, University of Padova, via Marzolo 5, Padova, Italy

^c Department of Chemistry and Biochemistry, The University of Texas at Arlington, Box 19065, Arlington, Texas 76019-0065, United States

Boron containing carbene complexes have never been investigated till now for their possible applications as anticancer drugs.¹ Our research group reported the synthesis of coinage metal-NHC complexes obtained from the precursors $\{[HB(RImH)_3]Br_2\}$ (R = Benzyl, Mesityl and t-Butyl)² and $\{[H_2B(BnTzH)_2]Br\}$.³

Here we present the synthesis of the new ligand $\{[H_2B(pNO_2BnTzH)_2]Br\}$ ³ and the related coinage carbene complexes. The synthesis of this ligand led to the isolation of the byproduct: $\{[HTz(pNO_2Bn)_2]Br\}$.⁴ This new ligand was also prepared by direct synthesis and used as carbene precursor for the formation of the related Cu(I), Ag(I), and Au(I) mono carbene complexes. Their cytotoxicity was evaluated over a panel of human cancerous cell lines and the results are here reported.




Figure 1.

References

- [1] Marinelli, M.; Santini, C.; Pellei, M., *Curr. Topics Med. Chem.*, submitted.
- [2] Biffis, A.; Gioia Lobbia, G.; Papini, G.; Pellei, M.; Santini, C.; Scattolin, E.; Tubaro, C., *J. Organomet. Chem.* **2008**, *693*, 3760-3766.
- [3] Papini, G.; Bandoli, G.; Dolmella, A.; Gioia Lobbia, G.; Pellei, M.; Santini, C., *Inorg. Chem. Commun.* **2008**, *11*, 1103-1106.
- [4] Marinelli, M., *et al.*, unpublished results.

Ru(II) and Cu(I) Complexes with Unsymmetrical N-Heterocyclic Carbenes Bearing N-Phenylpyrrole / N-Phenylindole Moiety

Krzysztof Grudzień,^{a*} Bartosz Trzaskowski,^b Karol Grela^a

^a Faculty of Chemistry, Biological and Chemical Research Centre, University of Warsaw, Żwirki i Wigury 101, 02-089 Warsaw, Poland.

Email: kgrudzien88@gmail.com

^b Centre of New Technologies, University of Warsaw, S. Banacha 2c, 02-097 Warsaw, Poland.

Exceptionally wide range of applications of n-heterocyclic carbenes (Figure 1) as organocatalysts and neutral ligands for transition metals comes from their easily tunable electronic and steric properties.¹ Our group specializes in olefin metathesis catalyzed by Ru(II) complexes with various ligands, including self-designed NHC's.² Recently we developed class of ruthenium catalysts with new, unsymmetrically substituted (R1, R2 = aryl, benzyl) NHC ligands (figure 2).^{3,4} Poster presentation will concern synthesis of their analogues bearing N-phenylpyrrole and N-phenylindole moieties, which were used in preparation of ruthenium(II) and copper(I) complexes.





Figure 1.



Abliialimov et. al., 2014

Figure 2.




Figure 3.

References

- [1] Hopkinson, M. N.; Richter, C.; Schedler, M.; Glorius, F. *Nature*, **2014**, *510*, 485-496.
- [2] Szczepaniak, G.; Kosiński, K.; Grela K., *Green Chem.*, **2014**, *16*, 4474-4492.
- [3] Abliialimov, O.; Kędziorek, M.; Torborg, C.; Malińska, M.; Woźniak, K.; Grela, K., *Organometallics*, **2012**, *31*, 7316-7319.
- [4] Abliialimov, O.; Kędziorek, M.; Malińska, M.; Woźniak, K.; Grela, K., *Organometallics*, **2014**, *33*, 2160-2171.

Ruthenium(II)-arene RAPTA Type Complexes Bearing Dibenzoylmethane: Organometallic Chemistry and Biological Activity

Agnese Petrini,^{a*} Francesca Condello,^a Jessica Palmucci,^b Giulio Lupidi,^a Massimo Nabissi,^a Antonio Rodríguez Diéguez,^c Riccardo Pettinari,^a Fabio Marchetti,^b Claudio Pettinari^a

^a School of Pharmacy and ^b School of Science and Technology, Università di Camerino, Via S. Agostino 1, 62032 Camerino (MC), Italy.
Email: agnese.petrini@unicam.it


^c Departamento de Química Inorgánica, Facultad de Ciencias, Universidad de Granada, 18071, Granada, Spain.

Half-sandwich η^6 -arene-ruthenium(II) complexes represent a very important class of organometallic compounds which have been extensively studied and appropriately modified to obtain compounds with various biological properties.¹

Dibenzoylmethane is a natural β -diketone that occurs in small amount in licorice roots. It exhibits several biological features such as activity against ischemic diseases² and different types of tumours.³

On the basis of our previous research on Ru(II)-arene complexes with acylpirazolones⁴ and curcuminoids⁵, in the present work we report the synthesis and the characterization of a series of arene-Ruthenium(II) (arene = *p*-cymene, hexamethylbenzene or benzene) complexes containing dibenzoylmethane and the ionic derivatives with PTA (1,3,5-triaza-7-phosphaadamantane) (Scheme 1).

Preliminary results on their biochemical studies and their antitumor activity evaluated *in vitro* in U266 and RPMI human multiple myeloma cell lines will also be reported.



Scheme 1.

References


- [1] Hartinger, C. G.; Metzler-Nolte, N.; Dyson, P. J. *Organometallics* **2012**, *31*, 5677.
- [2] Majeesh, N. J.; Willard, M. T.; Harris, W. B.; Sun, H.-Y.; Wang, R.; Zhong, H.; Umbreit, J. N.; Simons, J. W. *Biochem. Biophys. Res. Comm.* **2003**, *303*, 279.
- [3] a) Jackson, K. M.; DeLeon, M.; Reynold Verret, C.; Harris, Wayne B., *Cancer Letters* **2002**, *178*, 161; b) Weng, C.-J.; Yang, Y.-T.; Ho, C.-T.; Yen, G.-C., *J. Agric. Food Chem.* **2009**, *57*, 5235.
- [4] Marchetti, F.; Pettinari, R.; Pettinari, C. *Coordination Chemistry Reviews* **2015**, *303*, 1.
- [5] Pettinari, R.; Marchetti, F.; Condello, F.; Pettinari, C.; Lupidi, G.; Scopelliti, R.; Mukhopadhyay, S.; Riedel, T.; Dyson, P. J. *Organometallics* **2014**, *33*, 3709.

Syntheses of Ni and Au-N-Heterocyclic Carbene (NHC) Complexes and Their Catalytic Evaluation for C \equiv N and C \equiv C Activation

Kuldeep Singh and Jitendra K. Bera*

^a Department of Chemistry, Indian Institute of Technology Kanpur, Kanpur, 208016, India
E-mail: jbera@iitk.ac.in

The search for new atom-economical and green synthetic methods for the synthesis of functionalized important molecules has fascinated much attention owing to their potential industrial, medicinal and biological value.¹ Towards this goal new nickel and gold NHC complexes are synthesized and employed for amide synthesis by nitrile hydration and alkyne nitrogenation. Although a number of methods are known for the synthesis of amides, preparation under neutral conditions and without generating any waste is a challenging goal.² Nickel-NHC complex is very effective for hydration of nitriles under base free condition. Alkynes are successfully converted to amides by Au-NHC complex (Scheme 1). Industrially important acrylamide and antitubercular agent pyrazinamide are achieved in single step without producing any side product. Mechanisms of nitrile and alkyne activation will be discussed.



Scheme 1. Amide synthesis by Ni-NHC or Au-NHC catalyst.

References

- [1] (a) V. R. Pattabiraman, J. W. Bode, *Nature* **2011**, *480*, 471; (b) B. Shen, D. M. Makley, J. N. Johnston, *Nature* **2010**, *465*, 1027.
- [2] (a) P. Daw, A. Sinha, S. M. W. Rahaman, S. Dinda, J. K. Bera, *Organometallics* **2012**, *31*, 3790; (b) C. Qin, P. Feng, Y. Ou, T. Shen, T. Wang, N. Jiao, *Angew. Chem. Int. Ed.* **2013**, *52*, 7850.


Photocatalytic water oxidation with iridium catalysts under visible light

Ilaria Corbucci,^{a,*} Kevin Ellingwood,^b Pier Luigi Gentili,^a Alceo Macchioni^a

^a Department of Chemistry, Biology and Biotechnology, University of Perugia, Via Elce di Sotto 8, 06123, Perugia, Italy. *
ilariacorbucci@gmail.com

^b Montana State University, Bozeman, MT, 59717, United States.

The realization of an efficient apparatus for artificial photosynthesis, aimed at producing solar fuels, is strongly hampered by the difficulty of oxidizing water. Although several iridium complexes have demonstrated to be competent catalysts for water oxidation driven by chemical sacrificial oxidants,¹⁻⁶ such as cerium ammonium nitrate (CAN) or sodium periodate (NaIO₄), in only one case their photocatalytic activity has been documented.⁷ The classical protocol to explore possible photocatalytic activity stems on combining [Ru(bpy)₃]²⁺ as a photosensitizer and [S₂O₈]²⁻ as an electron acceptor as illustrated below.⁸⁻⁹



Scheme 1. Representation of a photoactivated cycle for water oxidation.

In this contribution, we report the results of a systematic study aimed at developing highly active molecular iridium photocatalysts and possibly correlating structure/performance using only the [Ru(bpy)₃]²⁺/[S₂O₈]²⁻ system. Generally speaking, all catalysts were found to be active and O₂ evolution occurred immediately after irradiation of the solution. Chemical and quantum yields varied in the ranges of 35-89%, based on the consumption of persulfate, and 7-22%, respectively, depending on the nature of ancillary ligands bonded at iridium.

References

- [1] McDaniel N. D., Coughlin, F.J., Tinker, L.L. and Bernhard, S.J. *J. Am. Chem. Soc.*, **2008**, *130*, 210.
- [2] Hull, J.F., Balcells, D., Blakemore, J.D. et al., *J. Am. Chem. Soc.*, **2009**, *131*, 8730.
- [3] Blakemore, J.D., Schley N.D., Balcells, D. et al., *J. Am. Chem. Soc.*, **2010**, *132*, 16017.
- [4] Savini, A., Bellachioma, G., Ciancaleoni, G. et al., *Chem. Commun.*, **2010**, *46*, 9218.
- [5] Dzik, W., Calvo, S.E., Reek, J.N.H. et al., *Organometallics*, **2011**, *30*, 372.
- [6] Bucci, A., Savini, A., Rocchigiani, L. et al., *Organometallics*, **2012**, *31*, 8071.
- [7] Volpe, A., Sartorel, A., Tubaro, C. et al., *Eur. J. Inorg. Chem.*, **2014**, *4*, 665.
- [8] Huang Z., Luo, Z., Geletii, Y.V., *J. Am. Chem. Soc.*, 2011, *133*, 2068.
- [9] Yamada Y., Yusuke Yamada, Y., Yano, K., Hong, D. and Fukuzumi, S., *Phys. Chem.* **2012**, *14*, 5753.

Tripodal Aminophenolate Iron(III) Complexes with Potential Biological Activity

Cristina P. Matos,^a M. Helena Garcia,^b João Costa Pessoa,^a Ana Isabel Tomaz,^b Isabel Correia^{a,*}

^a Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal. Email: icorreia@ist.utl.pt

^b Centro de Química Estrutural, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

The quest for cancer metallodrugs with improved pharmacological properties and different mechanisms of action than those of platinum drugs goes on. A possible approach to regulate the cytotoxic responses of metallodrugs is using biologically essential transition metals, such as iron, that plays an important role in many cellular processes. Iron, being redox active is involved in the regulation of cell-growth and differentiation.

Iron(III)-complexes of phenolate ligands with tripodal amines have attracted interest as mimics of enzyme active sites. The tetradentate ligands provide a reasonable analogue to histidine and tyrosinate coordination.¹ Substituents at the phenolate rings as well as the position and nature of the donor atoms are easily tunable features.² The introduction of a NN aromatic heterocyclic ligand in the coordination sphere could enforce the biological activity, as metal complexes containing phenanthroline derivatives are reported to be active against various pathologic conditions including cancer, microbial, and fungal infections.³ A wide array of electronic features and redox potentials are thus available, which can direct synthetic strategies towards attaining complexes with optimal performance. Tripodal aminophenolate iron(III) complexes have a well-developed chemistry, however their application as therapeutics for cancer is still an underdeveloped area of research open for further exploration. We report here the synthesis and preliminary studies on a family of iron(III) complexes composed of a [N,N-bis(2-hydroxy-3,5-dimethylbenzyl)-N-(2-pyridylmethyl)amine] ligand and different substituted phenanthrolines.




Figure 1.

References

- [1] Palaniandavar, M.; Velusamy, M.; Mayilmurugan, R. *J. Chem. Sci.* **2006**, *118*, 601.
- [2] Marshall-Roth, T.; Brown, S. N. *Dalton Trans.* **2015**, *44*, 677.
- [3] Jagadeesan, S.; Balasubramanian, V.; Baumann, P.; Neuburger, M.; Haussinger, D.; Palivan, C. G. *Inorg. Chem.*, **2013**, *52*, 12535.

Poster 47

Different Generations of Fluorescent PAMAM Dendrimers for Biomedical Applications

Cláudia S. Camacho, Dina Maciel,* Helena Tomás, João Rodrigues*

^a CQM- Centro de Química da Madeira, MMRG, Universidade da Madeira, Campus da Penteadá, 9000-390 Funchal, Portugal.

Email: joaor@uma.pt

The versatility of polyamidoamine (PAMAM) dendrimers has been demonstrated over these last years with applications in different areas, such as in the biomedical field¹ and, especially, as nanoplatforms for drug/gene delivery and bioimaging. However, these dendrimers are characterized by a weak intrinsic fluorescence which limits their application.¹ Even if fluorescence can be achieved by conjugation of labeling of fluorescent molecules (e.g. using fluorescein) with the peripheral functional groups of the dendrimer, the intrinsic fluorescent behavior of dendrimers can be further explored, being strongly influenced by the dendritic structure and/or the functional groups at the surface.¹⁻⁴ With the aim of exploring the fluorescence of dendrimers, in this work, generations 3 (G3) and 4 (G4) of amine-terminated PAMAM dendrimers were treated with ammonium persulfate (APS). The resulting compounds were characterized by different techniques, namely by Nuclear Magnetic Resonance, Ultraviolet-Visible, Fluorescence and Infrared Spectroscopies. Compounds presented a very intense fluorescence and absorption compared to pure PAMAM dendrimers. Particularly, solutions of the APS-treated dendrimer G3 and G4, when irradiated at 366nm, displayed an intense blue color luminescence (Figure 1). Additionally, preliminary results of the effect of different pHs over the intrinsic fluorescence and the stability of the APS treated dendrimers will be presented and discussed.




Figure 1. PAMAM G3.NH₂ dendrimer after oxidative treatment a) chemical structure b) irradiated under UV at 366nm.

Acknowledgments: This research was supported partially by Fundação para a Ciência e a Tecnologia (FCT) with Portuguese Government funds through the CQM Strategic Project PEst-OE/UI0674/2014 and the NMR Network – (PTNMR-2014 and PTNMR-2015). We acknowledge the continuous support of our work by Hotel Vidamar Resorts Madeira.

References

- [1] Tsai, Y.-J.; Hu, C.-C.; Chu, C.-C.; Imae, T. *Biomacromolecules*. **2011**, *12*, 4283.
- [2] Lee, W.I.; Bae, Y.; Bar, A.J. *J. Am. Chem. Soc.* **2004**, *126*, 8358.
- [3] Saravanan, G.; Abe, H. *J. Photochem. Photobiol., A* **2011**, *224*, 102.
- [4] Wang, D.; Imae, T. *J. Am. Chem. Soc.* **2004**, *126*, 1320.

Poster 48

Rational Improvement of Molecular Olefin Polymerization Catalysts

Francesco Zaccaria,^{a*} Christian Ehm,^a Peter H. M. Budzelaar,^b Vincenzo Busico^a

^a Dipartimento di Scienze Chimiche, Università di Napoli 'Federico II', Via Cintia, 80126 Napoli, Italy

^b Department of Chemistry, University of Manitoba, Winnipeg, MB R3T 2N2, Canada

The first challenge in polymer science is the controlled synthesis of macromolecules. In the case of polymers obtained by coordination catalysis, and specifically polyolefins, it is generally possible to tune polymer architecture and microstructure by modifying steric and (to some extent) electronic catalyst properties,¹ but this is still largely achieved by trial-and-error.

In this context, computational chemistry is mainly used to *interpret* – more than *predict* – structure/activity correlations. In the framework of a broader project on molecular design applications in polyolefin catalysis, we are currently investigating the factors which determine molar mass distributions of polymers produced with molecular catalysts, with special focus on electronic effects. A new computational strategy for *rational* average molar mass predictions is proposed. The approach is based on a detailed analysis of local minima "connected" to relevant transition states, and should represent a powerful tool to evaluate the balance between chain propagation and transfer by means of Density Functional Theory (DFT) calculations running in High Throughput Computation (HTC) mode.

To validate / benchmark it, we chose the industrially important class of Constrained Geometry Catalysts (CGC, Figure 1).² A number of precatalysts with different ligand substitution patterns have been synthesized, and their catalytic properties are being determined with High Throughput Experimentation (HTE) methods. The thus-obtained experimental database is being compared with the corresponding theoretical one. The first results of our study will be presented here.




Figure 1. Constrained Geometry Catalysts

References

- [1] Rappé, A. K.; Skiff, W.M.; Casewit, C. J. *Chem. Rev.* **2000**, *100*, 1435-1456
- [2] McKnight, A. L.; Waymouth, R. M. *Chem. Rev.* **1998**, *98*, 2587-2598

Poster 49

Modulation of the HCOOH / CO Ratio through the Use of Various Brønsted Acids for the Electroreduction of CO₂ Using [Mn(bpy-R)(CO)₃Br]-Type Catalysts

Luca Nencini,* Federico Franco, Claudio Cometto, Fabrizio Sordello, Claudia Barolo, Roberto Gobetto, Carlo Nervi, Claudio Minero

Dipartimento di chimica, Università di Torino, Via Pietro Giuria 7, 10125, Torino, Italia
e-mail: luca.nencini@unito.it

An efficient transformation of carbon dioxide into higher energy carbon products would make a remarkable impact on global economy, as an environmentally friendly decrease of the CO₂ content in the atmosphere could be coupled with a sustainable approach to gain useful chemicals (e.g. CO, HCOOH and CH₃OH).¹ In recent years, scientific research has primarily directed towards the discovery of new efficient organometallic molecular catalysts, containing abundant and cheap transition metals like Mn²⁺ and Group VI transition metals.³

Taking inspiration from Saveant's⁴ work where in the Fe(III) porphyrins, availability of local proton sources is known to greatly enhance the selectivity and the redox catalytic activity for CO₂ reduction to CO, we designed some tricarbonyl Manganese coordination compounds containing polypyridyl ligands⁵, [Mn(bpy-R)(CO)₃Br] (bpy-R = 4-phenyl-6-(1,3-dihydroxybenzen-2-yl) 2,2'-bipyridine (**1**) and 4-phenyl-6-(1,3-dimethoxybenzen-2-yl) 2,2'-bipyridine (**2**)). Our studies showed that the electrochemical of a novel polypyridyl Mn(I) catalyst containing two acidic OH groups in proximity of the purported metal binding site for CO₂ redox catalysis is reported. **1** showed a substantial catalytic activity in anhydrous media induced by the presence of a local proton source. The electrochemical data for **1** highlighted two crucial issues: (1) CV exhibited a strong catalytic effect on the current of the 1/2' reduction when the solution was saturated with CO₂, so that the 2e⁻ pathway was supposed to be favoured for **1**. This represents the first reported experimental evidence of an intramolecular proton-assisted catalytic process for a Mn(I) catalyst. (2) Bulk electrolysis at the end of the catalytic plateau (1.8 V) under CO₂ showed an unusual change in selectivity for CO₂ reduction by the Mn(I) catalyst, giving a mixture of CO and HCOOH. The catalytic mechanism is under investigation by techniques of IR spectroelectrochemistry and DFT calculations.

References

- [1] (a) S. Berardi, S. Drouet, L. Francas, C. Gimbert-Surinach, M. Guttentag, C. Richmond, T. Stolla and A. Llobet, *Chem. Soc. Rev.*, **2014**, 43, 7501-7519; (b) J. Qiao, Y. Liu, F. Hong and J. Zhang, *Chem. Soc. Rev.*, **2014**, 43, 631-675; (c) A. Taheri Najafabadi, *Int. J. Energy Res.*, **2013**, 37, 485-499.
- [2] Bourrez, M. et al., *Angew Chem*, **2011**, 50, 9903-9906.
- [3] Nencini, L.; Franco, F.; Cometto, C.; Nervi, C.; Gobetto, R.; Sordello, F.; Minero, C. *ChemElectroChem*, **2015**,
- [4] C. Costentin, S. Drouet, M. Robert and J. M. Saveant, *Science*, **2012**, 338, 90-94.
- [5] Franco, F.; Cometto, C.; Nervi, C.; Gobetto, R.; Sordello, F.; Minero, C.; Vallana, F.; Priola, E., *Chem Comm* **2014**, 50, 14670-14673.

Poster 50

Synthesis of Diketopyrrolo[3,4-c]pyrrole -Based Derivatives for Organic Electronics

Angela Punzi,^a Cosimo V. Ciasca,^{a*} Eliana Maiorano,^a Francesca Nicoletta,^b Gianluca M. Farinola.^{a*}

^a Università degli studi di Bari "Aldo Moro", Via Orabona 4, 70125 Bari, Italy, cosimo.ciasca@uniba.it

^b Università degli studi di Catania, Piazza Università 2, 95131 Catania, Italy

Stereoselective In recent years, diketopyrrolo[3,4-c]pyrrole (DPP) moiety has been widely used as a building block in the synthesis of a variety of organic semiconductor materials for device applications, particularly for organic thin film transistors (OTFTs) and organic photovoltaics (OPVs).¹ Among all, the thiophene-functionalized DPP moiety TDPP has attracted extensive attention because of its unique properties arising from strong intramolecular charge transfer and good π - π stacking of the resulting conjugating units. A strong influence on these properties is also due to N-substituents of the lactam rings as well as the side chains on conjugate backbone. Recently, interesting materials for applications in organic electronics have been obtained by introduction of hydrophilic oligoether chains as stack-inducing agents on the TDPP units. More relevant, aromatic end-groups appended to the central TDPP core of symmetrically substituted small molecules, typically introduced by cross-coupling reactions,² strongly impact on the efficiency of these molecules in BHJ devices. We report the synthesis of new TDPP-based molecular semiconductors in which 1,2,3-triazole rings are conjugated to the central TDPP core (Figure1). The 1,2,3-triazoles are aromatic heterocycles with high stability which are capable of active participation in hydrogen bonding as well as dipole-dipole and π - π stacking interactions.³ A variable number of hydrophilic oligoether chains were introduced on the innovative conjugate backbone in order to achieve compounds with a tailored solubility in environmental friendly polar solvents, such as alcohols or water, potentially useful for green OPV or biological applications.




Figure 1.

References

- [1] Nielsen, C. B.; Turbiez, M.; McCulloch I. *Adv. Mater.* **2013**, 25, 1859.
- [2] Marzano, G.; Ciasca, C. V.; Babudri, F.; Bianchi, G.; Pellegrino, A.; Po, R.; Farinola, G. M. *Eur. J. Org. Chem.* **2014**, 30, 6583.
- [3] Whiting, M.; Muldoon, J.; Lin, Y.-C.; Silverman, S. M.; Lindstrom, W.; Olson, A. J.; Kolb, H. C.; Finn, M. G.; Sharpless, K.B.; Elder, J. H.; Fokin V. V. *Angew. Chem. Int. Ed.* **2006**, 45, 1435.


Revisiting Ammonium Formate: Chemoselective Transfer Hydrogenation of Aldehydes Catalyzed by Ruthenium Pincer Complexes

Salvatore Baldino,^{a*} Walter Baratta,^{a*} Steven Giboulot,^a Sarah Facchetti,^b Antonio Zanotti-Gerosa,^b Hans G. Nedden^b

^a Dipartimento di Chimica, Fisica Ambiente, Università di Udine, Via Cotonificio 108, I-33100 Udine, Italy *salvatore.baldino@uniud.it; walter.baratta@uniud.it

^b Johnson Matthey Catalysis and Chiral Technologies, 28 Cambridge Science Park, Milton Road, Cambridge, CB4 0FP, United Kingdom

Bis(pyrazolyl)alkanes Ammonium formate (AF) is a cheap, readily accessible and versatile reagent which has been largely employed across a wide range of organic transformations.¹ In the last decade AF has been successfully used in transition-metal-catalyzed reduction processes, particularly in the reductive amination of ketones,² given its dual nature of both hydrogen and ammonia donor. Amongst late transition metal compounds, ruthenium complexes have emerged as very productive catalysts for hydrogenative reactions of carbonyl substrates.³ We have recently studied the catalytic properties of RuCl(CNN^{Ph})(PP) complexes **1-3** in the transfer hydrogenation of aldehydes using 2-propanol as hydrogen source.⁴ Then we extended the application of RuCl(CNN^{Ph})(PP) compounds in catalytic hydrogenative reactions mediated by different hydrogen sources and we report here the unprecedented use of AF in the chemoselective and highly productive reduction of commercial-grade aldehydes to their corresponding primary alcohols catalyzed by ruthenium pincer complexes **1-3**.



References

- [1] Ram, S.; Spicer, L. D.; *Tetrahedron Lett.* **1988**, *29*, 3741.
- [2] Kadyrov, R.; Riermeier, T. H.; *Angew. Chemie Int. Ed.* **2003**, *115*, 5630.
- [3] Chelucci G.; Baldino S.; Baratta W.; *Coord. Chem. Rev.* **2015**, *300*, 29.
- [4] Unpublished results.

DNA, BSA Binding and Cytotoxic Properties of Iron(III) Complex Bearing Formazan Ligands

Nuno M. R. Martins,^{a*} Sellamuthu Anbu,^a Kamran T. Mahmudov,^{a,b} Rajendran Ravishankaran,^c M. Fátima C. Guedes da Silva,^a Luísa M. D. R. S. Martins,^{a,d} Anjali A. Karande,^c Armando J. L. Pombeiro^a

^a Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisbon, Portugal. E-mail: nunomartins@tecnico.ulisboa.pt ^b Department of Chemistry, Baku State University, Baku, Azerbaijan; ^c Department of Biochemistry, Indian Institute of Science, Bangalore-560 012, India; ^d Chemical Engineering Department, ISEL, R. Conselheiro Emídio Navarro, Lisboa, Portugal.

Cytotoxicity of most of the metal complexes has been mainly correlated with their ability to bind and damage DNA or protein.^[1] Among them, as alternative to platinum-based drugs, iron(III) complexes have emerged as one of the leading players in this field.^[2] However, DNA and protein interacting studies as well as cytotoxic properties of water soluble iron(III) complexes derived from formazan ligands have not yet been reported. Herein we report the synthesis of the Fe^{III} complex [Fe(κ^3 -HL)₂] (**1**) that was isolated by treatment of iron(III) chloride hexahydrate with the new (1*E*,1*E*)-*N*'-2-di(1*H*-1,2,4-triazol-3-yl)diazene-carbohydrazonyl cyanide (**H₃L**), and characterized by elemental analysis, IR spectroscopy and single crystal X-ray diffraction. The interaction of calf thymus DNA (CT DNA) and bovine serum albumin (BSA protein) with complex **1** has been investigated by absorption, fluorescence spectroscopic and viscosity measurement techniques. The observed intrinsic DNA binding constant ($1.98(\pm 0.26) \times 10^5 \text{ M}^{-1}$), number of binding sites ($s \leq 1$) of complexes to DNA and viscosity data suggest groove binding and/or a partial intercalative mode of binding to CT DNA. Complex **1** shows good binding propensity to the BSA, giving K_{BSA} values of $1.06(\pm 0.08) \times 10^6 \text{ M}^{-1}$. Complex **1** was evaluated for its *in vitro* cytotoxic activity against the human cervical (HeLa) and breast (MCF-7) cancer cell lines as well as noncancer breast epithelial (MCF-10a) cell lines. The results showed that **1** mediates a more strong cytotoxic response to the tested cancer cell lines than normal ones.




Figure 1. Synthesis of a new Fe^{III} complex with heterocyclic formazan (**1**) which shows DNA, BSA binding and anticancer abilities.

Acknowledgements: This work has been partially supported by the FCT, Portugal (UID/QUI/00100/2013). N.M.R.M and S.A. are thankful to FCT for the PhD (SFRH/BD/52371/2013) and post-doc fellowship (SFRH/BPD/76451/2011). R.R. is thankful to the CSIR-SRF, India. The authors acknowledge the Portuguese NMR Network (IST-UTL Centre) for access to the NMR facility, and the IST Node of the Portuguese Network of mass-spectrometry for the ESI-MS measurements.

References

- [1] a) S. Anbu, S. Kamalraj, B. Varghese, J. Muthumary, M. Kandaswamy, *Inorg. Chem.* **2012**, *51*, 5580–5592; b) S. Anbu, A. Paul, R. Ravishankaran, M. F. C. Guedes da Silva, A. A. Karande, A.J.L. Pombeiro, *Inorg. Chim. Acta* **2014**, *423*, 183–193; c) K.Saleem, W. A Wani, A. Haque, M. N. Lone, M.-F. Hsieh, M. A. Jairajpuri, I. Ali, *Fut. Med. Chem.* **2013**, *5*, 135–146; d) M. Ebrahimi, T. Khayamian, H. Hadadzadeh, B. E. S. Tabatabaei, Z. Jannesari, G. Khaksar, *J. Biomol. Struct. Dyn.* **2015**, *11*, 1–14.
- [2] a) B.-L. Fei, W.-S. Xu, W.-L. Gao, J. Zhang, Y. Zhao, J.-Y. Long, C. E. Anson, A. K. Powell, *J. Photoch. Photobio. B* **2014**, *142C*, 77–85; b) Y.-N. Shi, K. Zheng, L. Zhu, Y.-T. Li, Z.-Y. Wu, C.-W. Yan, *J. Biochem. Mol. Toxicol.* **2015**, *29*, 221–233.

Poster 53

Novel Hoveyda Type Metathesis Catalysts Bearing Sterically Hindered, Unsymmetrical NHC Ligand

Paweł Malecki,^a Karol Grela^a

^a Biological and Chemical Research Center, Faculty of Chemistry, University of Warsaw, Żwirki i Wigury 101, 02 089 Warsaw, Poland.
Email: pawelmalecki.chem@gmail.com, karol.grela@gmail.com

In 2000 A. Hoveyda developed a new type of olefin metathesis ruthenium catalyst, commonly referred as Hoveyda-Grubbs catalyst. Since then a lot of structure modifications based on this motif were made.¹

However finding an ideal catalyst that is both active and stable is still a challenge. One of the most promising modifications is the change in NHC ligand. Our group follows that trend and focuses on synthesizing and modifying those types of ligands.²

We synthesized a variety of new Hoveyda type complexes possessing bulky unsymmetrical N-heterocyclic carbene (NHC) (fig.1). Those complexes are potentially interesting in cross metathesis reaction, especially when one of the olefins is ethylene. During the presentation the synthesis and reactivity of the new complexes will be discussed.




Figure 1.

References

- [1] Vougioukalakis, G.C.; Grubbs, R.H. *Chem Rev* **2010**, *110*, 1746
[2] Abllialimov, O.; Kędziołek, M.; Malińska, M. Wozniak, K.; Grela, K. *Organometallics*, **2014**, *33*, 2160.

Poster 54

Comparing the Catalytic Activity of Cp*Ir Catalysts with bpy and dpa: Insights into the Mechanism of Water Oxidation Driven by CAN

Alberto Bucci,^{a*} Gabriel Menendez Rodriguez,^a Gianfranco Bellachioma,^a Cristiano Zuccaccia,^a Alceo Macchioni^a

^a Department of Chemistry, Biology and Biotechnology, Via Elce di Sotto 8, 06123 Perugia (PG), Italy.
Email: albechm@gmail.com

The rapid depletion of fossil fuels asks for the utilization of alternative and sustainable energy sources. Among them, the sunlight has attracted great attention especially as for the realization of a photosynthetic apparatus aimed at producing solar fuels, exploiting the electrons coming from the catalytic oxidation of water to O₂.¹ Unfortunately, the performances of water oxidation catalysts (WOCs) reported in the literature², are far from those required for real applications.

Cp*Ir complexes (Cp* = 1,2,3,4,5-pentamethylcyclopentadiene) are highly active and robust WOCs and their performances can be easily tuned by the choice of the ancillary ligand. Herein we report the study of the catalytic behavior of [Cp*Ir(dpa)Cl₂] (dpa = 2,2' dipyridilamine) (**1**) using CAN (cerium ammonium nitrate) as sacrificial oxidant. The performances of **1** were strictly compared to those of its analogous [Cp*Ir(bpy)Cl₂] (**2**) previously reported by us^{3,4}. The presence of the -NH bridge substantially increases the activity of **1** (up to 6 times) if compared to that of **2**; this led us to carry on deeper kinetic studies, in order to rationalize such difference. Those studies shed some new insights on the reaction mechanism of water oxidation driven by CAN in general. Indeed, they clearly showed that CAN is consumed before the oxygen evolution starts (Figure 1), thus suggesting a mechanism proceeding through the formation of an intermediate species that maintains the oxidative power of Ce⁴⁺ or traps the preformed O₂. We believe that such a mechanism might be quite general, since similar results were observed for other molecular catalysts and also for IrO₂.




Figure 1.

References

- [1] Balzani, V.; Credi, A.; Venturi, M. *ChemSusChem*, **2008**, *1*, 26–58
[2] Blackmore, J. D.; Crabtree, R.H.; Brudvig, W. G. *Chem. Rev.* DOI: 10.1021/acs.chemrev.5b00122
[3] Savini, A.; Bellachioma, G.; Ciancaleoni, G.; Zuccaccia, C.; Zuccaccia, D.; Macchioni, A. *Chem. Commun.* **2010**, *46*, 9218-9219.
[4] Savini, A.; Bucci, A.; Bellachioma, G.; Rocchigiani, L.; Zuccaccia, C.; Llobet, A.; Macchioni, A. *Eur. J. Inorg. Chem.* **2014**, 690–697.

Silylated and Germylated Pyrazoleboronic Acids: Synthesis and Characterization

Agnieszka Górska, Tomasz Kliś, Janusz Serwatowski


Faculty of Chemistry, Warsaw University of Technology, Poland.

Email: agorska@ch.pw.edu.pl

Cross-coupling reactions of organoboron compounds discovered by A. Suzuki are important in various fields of organic synthesis and combinatorial chemistry. Among a relatively wide assortment of available aryl- and heteroarylboronic acids used for Suzuki reaction, the derivatives of pyrazole are still rare species. On the other hand, the pyrazole fragment is present in a large number of natural and synthetic small ligands that interact with various enzymes and receptors of pharmacological significance.

5-Membered nitrogen-containing heteroaromatics (azoles) can often be functionalized efficiently by lithiation of different ring positions and exocyclic α -position. In this work reactions of many 4-bromopyrazole derivatives with LDA were investigated. Obtained lithium derivatives were reacted with various electrophiles as: Chlorotrimethylgermane, Chlorotrimethylsilane or Chlorodimethylsilane to form the corresponding substituted compounds. These products were treated as intermediates for the following reactions with *t*-BuLi where the bromine was substituted (or not) with lithium. Those which underwent exchange reaction were reacted with $B(OEt)_3$ and hydrolyzed to form finally the corresponding boronic acids.

The structure of diboronic acids was studied using multinuclear NMR spectroscopy and elemental analysis. The crystalline derivatives obtained were suitable for X-ray measurements.



References

- [1] Ivachtchenko, A. V.; Kravchenko, D. V.; Zheludeva, V. Y.; Pershin, D. G. *J. Heterocyclic Chem.*, **2004**, 931.
- [2] Balle, T.; Begtrup, M.; Jaroszewski, J. W.; Liljefors, T.; Norrby P. *Org. Biomol. Chem.*, **2006**, 4, 1261.
- [3] Durka, K.; Górska, A.; Kliś, T.; Kublicki, M.; Serwatowski, J.; Woźniak, K. *Tetrahedron Lett.*, **2015**, 56, 1855-1859.

High Performance Isocyanide Scavengers for Use in Low-Waste Purification of Olefin Metathesis Products

Grzegorz Szczepaniak,^{a*} Katarzyna Urbaniak,^b Celina Wierzbicka,^b Krzysztof Kosiński,^a Krzysztof Skowerski,^b Karol Grela^{a*}

^a Biological and Chemical Research Centre, Faculty of Chemistry, University of Warsaw, Żwirki i Wigury 101, 02-089 Warsaw, Poland

Email: szczepaniak.grzegorz@gmail.com ; karol.grela@gmail.com

^b Apeiron Synthesis, Duńska 9, 54-427 Wrocław, Poland.

In recent years, more attention is focused on tackling the problem of metal contamination as early as possible in the synthetic pipeline.¹ Possible approaches include heterogeneous catalysis, self-scavenging catalysts, and the use of metal scavengers.² Recently, we reported an olefin metathesis catalyst with quaternary ammonium groups in its NHC ligand, which could be easily removed from the solution by filtration through a short plug of silica gel or extraction.³

In this work three isocyanides (**I-III**) containing tertiary nitrogen atoms and between one and three isocyanide groups were used as metal scavengers (Figure 1).⁴ These compounds prove to be highly effective at removing ruthenium residues from post-reaction mixtures, in many cases reducing the ruthenium content below 0.0015 ppm and requiring short binding times and low stoichiometric excesses. Finally, we report on the synergy between the scavenger **II** and one of the previously reported self-scavenging catalysts, which allows one to obtain highly pure products even in highly challenging cases. Additionally, isocyanide **II** is shown to be a quenching agent for olefin metathesis, superior to commonly used ethyl vinyl ether.




Figure 1

References

- [1] Vougioukalakis, G. C. *Chem. Eur. J.* **2012**, 18, 8868.
- [2] Szczepaniak, G.; Kosiński, K.; Grela, K. *Green Chem.* **2014**, 16, 4474.
- [3] Skowerski, K.; Wierzbicka, C.; Szczepaniak, G.; Gulajski, L.; Bieniek, M.; Grela, K. *Green Chem.* **2012**, 14, 3264.
- [4] Szczepaniak, G.; Urbaniak, Wierzbicka, C.; K.; Kosiński, K.; Skowerski, Grela, K. *submitted*.

Poster 57

Synthesis of Thiophene Derivatives in DES

Raffaella Mancuso,^{a,b} Asif Maner,^{a*} Vito Capriati,^c Bartolo Gabriele^a

^aDipartimento di Chimica e Tecnologie Chimiche, Università della Calabria, Via Pietro Bucci, 12/C, 87036 Arcavacata di Rende (CS), Italy.


Email: asif.maner@unical.it

^bDipartimento di Ingegneria Meccanica, Energetica e Gestionale, Università della Calabria, 87036 Arcavacata di Rende (CS), Italy


^cDipartimento di Farmacia-Scienze Del Farmaco, Università di Bari Aldo Moro, Consorzio C.I.N.M.P.I.S., Via E. Orabona 4, Bari, Italy

Deep eutectic solvents (DESs), which can be defined as molecular complexes typically formed between a simple halide salt (i.e. choline chloride) and a hydrogen bond donor, have found a wide variety of applications in different fields of modern chemistry, including organic synthesis, biocatalytic reactions, dissolution of metal oxides, electrodeposition of metals, and materials chemistry. However the number of studies reporting the ability of DESs to serve as green and biorenewable reaction media in metal-catalysed organic reactions is still scarce and no examples of iodocyclization reactions in DESs are reported in literature.

We report here the first example of iodocyclisation reaction in DES. Readily available 1-mercapto-3-yn-2-ols **1** are converted into the corresponding 3-iodothiophenes **2** under base-free conditions, at room temperature and in Choline Chloride/Glycerol (1*ChCl*/2*Gly*) such as solvent (eq.1). Products **2** are obtained in good yields (60-79%) and the DES can be recycled different times without loss of activity.



Starting from 1-mercapto-3-yn-2-ols **1** we have also obtained thiophenes **3** by PdI₂-catalyzed heterocyclization reaction (eq. 2). The reaction is carried out under air, at room temperature and in 1*ChCl*/2*Gly* such as solvent. Thiophenes **3** are formed in good to high yields (63-81%) and with the possibility to recycle DES-catalyst system several times without affecting the reaction.



Poster 58

Manganese(II) Complexes with Pyrazole Ligands: Synthesis and Oxidative Catalysis

Tiago A.G. Duarte,^{a*} Luísa M.D.R.S. Martins^{a,b} Ana P. Carvalho,^c Armando J.L. Pombeiro^a

^a Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001

Lisboa, Portugal.

Email: tiago.gomes.duarte@tecnico.ulisboa.pt


^b Chemical Engineering Department, Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa, R.

Conselheiro Emídio Navarro, 1959-007 Lisboa, Portugal.

^c Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016

Lisboa, Portugal.

Oxidation reactions catalyzed by transition metal complexes are quantitatively the most important homogeneously catalyzed reactions in chemical industry. The use of metallic complexes with C homoscorpionate poly(pyrazol-1-yl)methane ligands as oxidation catalysts is experiencing significant development, mainly due to their coordinative versatility that allows the electronic and steric features at a metal center to be tuned.^{1,2} Herein we report the design of manganese(II) complexes containing the C-homoscorpionate tris(pyrazol-1-yl)methane ligand, HC(pz)₃ (pz = pyrazolyl), or pyrazole (Hpz), formed by the reactions depicted in Scheme 1, for application as catalysts for the partial oxidation of saturated hydrocarbons, a field of great interest due to the considerable inertness of the substrates. The effects of some catalytic parameters on the oxidation reactions are also discussed.



Scheme 1.

Acknowledgements: This work has been partially supported by the Foundation for Science and Technology (FCT), Portugal (UID/QUI/00100/2013 and UID/MULTI/00612/2013). Tiago A.G. Duarte is thankful to FCT for his PhD fellowship (PD/BD/105993/2014). The authors acknowledge the Portuguese NMR Network (IST-UTL Centre) for access to the NMR facility, and the IST Node of the Portuguese Network of mass-spectrometry for the ESI-MS measurements

References

- [1] Silva, T. F. S., Martins, L. M. D. R. S., M. Guedes da Silva, M. F. C., Kuznetsov, M. L., Fernandes, A. R., Silva, A., Pan, C.-J., Lee, J.-F., Hwang, B.-J., Pombeiro, A. J. L., *Chem. Asian J.* **2014**, *9*, 1132.
- [2] Martins, L. M. D. R. S., Pombeiro, A. J. L., *Coord. Chem. Rev.* **2014**, *265*, 74.

Straightforward assembly of bis(benzoxaborole) systems from 2-formylphenylboronic acids

Krzysztof M. Borys,^a Alicja Matuszewska, Agnieszka Adamczyk-Woźniak, Andrzej Sporzyński

Faculty of Chemistry, Warsaw University of Technology, Noakowskiego 3, 00-664 Warsaw, Poland, *k.borys@ch.pw.edu.pl

Benzoxaboroles are cyclic, internal hemiesters of phenylboronic acids. These organoboron compounds have found wide applications, ranging from medicine and biological chemistry to materials chemistry and organic synthesis.^{1,2} Due to a growing interest in diversely functionalized benzoxaboroles, novel and effective methods for their preparation are highly desirable.

The developed methodology enables a straightforward and efficient synthesis of piperazine-based bis(benzoxaboroles) (Figure 1).^{3,4} The method originates from the studies on the reactivity of 2-formylphenylboronic acid towards secondary amines.⁵ Herein, the product is formed spontaneously in ethereal solution *via* dehydrative amination of 2-formylphenylboronic acid (or its derivative) with piperazine.

Interestingly, bis(benzoxaborole) **1** was found to have considerable biological activity, being an effective antifungal agent.⁶




Figure 1.

Acknowledgements: Krzysztof M. Borys acknowledges the Ministry of Science and Higher Education of Poland for financial support within the "Diamantowy Grant" program (grant no. DI 2012 015042 (0150/DIA/2013/42)), as well as the 10th ISOC Organizing Committee for the conference grant.

References

- [1] Adamczyk-Woźniak, A.; Borys, K. M.; Sporzyński, A. *Chem. Rev.* **2015**, *115*, 5224.
- [2] Liu, C. T.; Tomsho, J. W.; Benkovic, S. J. *Bioorg. Med. Chem.* **2014**, *22*, 4462.
- [3] Adamczyk-Woźniak, A.; Borys, K. M.; Madura, I. D.; Michalek, S.; Pawelko, A. *Tetrahedron* **2013**, *69*, 8936.
- [4] Adamczyk-Woźniak, A.; Ejsmont, K.; Gierczyk, B.; Kaczorowska, E.; Matuszewska, A.; Schroeder, G.; Sporzyński, A.; Zarychta, B. *J. Organomet. Chem.* **2015**, *788*, 36.
- [5] Adamczyk-Woźniak, A.; Madura, I.; Velders, A. H.; Sporzyński, A. *Tetrahedron Lett.* **2010**, *51*, 6181.
- [6] Wieczorek, D.; Lipok, J.; Borys, K. M.; Adamczyk-Woźniak, A.; Sporzyński, A. *Appl. Organometal. Chem.* **2014**, *28*, 347.

Heck arylation of allyl alcohol catalyzed by palladium compounds

Stanisława Tarnowicz,^{a*} Waleed Alsalahi,^a Ewa Mieczysława,^a Anna M. Trzeciak^a

^a Faculty of Chemistry, University of Wrocław, 14 F. Joliot-Curie St., 50-383 Wrocław, Poland.

Email: stanislaw.tarnowicz@chem.uni.wroc.pl

Palladium-catalyzed Heck reaction with allylic alcohols as substrates is interesting because it generates variety of arylated ketones or aldehydes as products. β -Aryl ketones, products of this type Heck reaction can be used for the synthesis of medicinal products, for example 4-phenyl-2-butanones (enzymatic inhibitors).¹ Product A (3-phenylpropionaldehyde) might have synergize effect with trans-cinnamaldehyde for antibacterial and anti-inflammatory effects on *H. pylori* infection.²




Figure 1. Heck arylation of allyl alcohol catalyzed by palladium catalyst.

In this work we would like to present our result of synthesis of arylated ketones, involving different aryl halides and catalysed by Pd(OAc)₂, PdCl₂·COD or Pd(0) nanoparticles. The reaction was studied with variation of parameters such as different temperature (85°C - 130°C), time (1.5 h – 24 h) and addition of molten salts (TBAB). We have obtained aldehydes A and B as the main products and they were isolated using extraction and column chromatography on silica gel. The best system for iodobenzene contained NaOAc as base. In 85°C, after 1.5 h, we obtained 79% product A, 13 % B and 2% C. The presence of salt (TBAB) increased yield (49%) of the main product (A) when PhBr was used as aryl halide. We received product A with the yield 48-79% in reaction with para-substituted aryl iodides and the best result were 79% for 4-Iodoanisole and 71% for 4-Iodophenol.

References

- [1] Dearden, J. C.; Nicholson, R. M. *J. Pharm. Pharmacol.* **1984**, *36*, 713.
- [2] Fang, S. B.; Ko, H. Y.; Huang, S. T.; Huang, C. H.; Li, L. T.; Chen, C. C.; Wang, K. C.; Pai, C. P.; Lee, H. C.; Fang, H. W. *RSC Adv.* **2015**, *5*, 22097.

Poster 61

Formic Acid as an Alternative Hydrogen Source in Ruthenium Catalyzed Transfer Hydrogenation of Olefins

Grzegorz K. Zieliński,^{a,*} Cezary Samojłowicz,^a Tomasz Wdowik,^a Karol Grela,^{a,b}

^a Institute of Organic Chemistry PAS, Kasprzaka 44/52, 01-224, Warsaw, Poland

^b Biological and Chemical Research Centre, Faculty of Chemistry, University of Warsaw, Żwirki i Wigury 101, 02-089, Warsaw, Poland; karol.grela@gmail.com

Catalytic olefin metathesis, as one of the most important methodologies of C-C double bond formation, does not need a particular introduction. Furthermore ruthenium alkylidene complexes were found not only as a great olefin metathesis catalysts¹ but they are well known in literature as active catalysts of numerous non-metathetic reactions.²

During our recent study on application of a new nitronate ruthenium complex **1** in olefin metathesis, we have noticed its unexpected non-metathetic activity. Depending on the conditions applied, the same complex promoted efficiently olefin metathesis, cycloisomerization, reduction of a carbonyl group, as well as isomerization of a C-C double bond.³

The interesting further results confirmed that complex **1** can also catalyze metathesis and transfer hydrogenation in one-pot sequence without isolation of the metathesis product. Moreover we proved that in this conditions many commercially available ruthenium alkylidene complexes (like **Gru-II**) can catalyze metathesis and transfer hydrogenation sequence or transfer hydrogenation depending on used substrate (**Figure 1**). Methodology was tested on several olefins and more interesting results appeared. The methodology that was developed in our group allows for selective hydrogenation in mild conditions using formic acid instead of dangerous hydrogen gas and does not cleave the benzyl groups.⁴




Figure 1.

References

- [1] Grela, K., Ed. *Olefin Metathesis: Theory and Practice*, John Wiley & Sons, Inc., Hoboken, New Jersey, **2014**.
- [2] Alcaide, B.; Almendros, P.; Luna, A. *Chem. Rev.* **2009**, *109*, 3817.
- [3] a) Wdowik, T.; Samojłowicz, C.; Jawiczuk, M.; Grela, K. Polish Patent Application, P.396439, **2011**; b) Wdowik, T.; Samojłowicz, C.; Jawiczuk, M.; Grela, K. International Patent Application, PCT/IB2012/055058, **2012**; c) Wdowik T.; Samojłowicz C.; Jawiczuk M.; Malińska M.; Woźniak K.; Grela K. *Chem. Commun.* **2013**, *49*, 674.
- [4] Zieliński G. K.; Samojłowicz, C.; Wdowik, T.; Grela, K. *Org. Biomol. Chem.*, **2015**, *13*, 2684.

Poster 62


CO₂ Reduction to Methanol and Methylamines Using Formic Acid as a C–H Bond Shuttle

Solène Savourey,^{*} Guillaume Lefèvre, Jean-Claude Berthet, Pierre Thuéry, Caroline Genre and Thibault Cantat

^a LCMCE, CEA Saclay, 91191 Gif-sur-Yvette cedex.

Email: solene.savourey@cea.fr

Efficient conversion of CO₂ to methanol is a key process to reach a methanol economy, based on a closed carbon cycle.¹ Such goal could be achieved by the 6-electron reduction of CO₂ or its hydrogenation to methanol. However both solutions currently suffer from low faradaic efficiencies. An interesting alternative would consist in utilizing formic acid (FA) as a C–H bond shuttle in the reduction of CO₂ to methanol. This strategy relies on the 2–electron reduction of CO₂ to FA, in an electrochemical cell, and this methodology is now technically and economically available, thanks to efficient electrocatalysts.² Disproportionation of FA is then required to produce methanol. Miller *et al.* showed, for the first time in 2013, that a molecular complex could promote the disproportionation of FA to methanol.³ Though promising, this strategy currently suffers from the use of expensive iridium catalysts and the yields of methanol do not exceed 1.9 %.³ We reported the efficient disproportionation of FA to methanol, with methanol yields of up to 50.2 %, using ruthenium molecular catalysts. Different pathways involving transient ruthenium–hydride species have been unveiled, based on mechanistic experimental and DFT investigations.⁴



The utilization of formic acid as a C-H bond shuttle in the conversion of CO₂ can also be extended to the synthesis of fine chemicals. In this respect, the use of CO₂ as a carbon source for the methylamine has been recently unveiled and this methodology requires the use of external reductants such as hydrosilanes and molecular hydrogen.⁵ To circumvent the use of additional reductants, we have recently demonstrated that formic acid could be used as the unique source of carbon and hydrogen in the catalytic methylation of amines.⁶

References


- [1] G. Olah, A. Goepfert, G. K. Surya Prakash, *Beyond Oil and Gas: The Methanol Economy*, Wiley- VCH, Weinheim, **2009**.
- [2] H.-R. Jhong, S. Ma, P. J. A. Kenis, *Curr. Opin. Chem. Eng.* **2013**, *2*, 191.
- [3] A. J. M. Miller, D. M. Heinekey, J. M. Mayer, K. I. Goldberg, *Angew. Chem. Int. Ed.* **2013**, *52*, 3981; *Angew. Chem.* **2013**, *125*, 4073.
- [4] S. Savourey, G. Lefèvre, J.-C. Berthet, P. Thuéry, C. Genre, T. Cantat, *Angew. Chem., Int. Ed.*, **2014**, *53*, 10466.
- [5] A. Tlili, E. Blondiaux, X. Frogneux, T. Cantat, *Green. Chem.* **2014**, *10*, 1039/C4GC01614A.
- [6] S. Savourey, G. Lefèvre, J.-C. Berthet, T. Cantat *Chem. Commun.*, **2014**, *50*, 14033.+

Zn(II) Homogeneous Supported Catalysts for the Efficient Esterification and Transesterification of Vegetable Oils

Vincenzo Benessere, Maria E. Cucciolito, Martino Di Serio, Roberto Esposito,* Veronica Papa, Francesco Ruffo, Rosa Turco, Rosa Vitiello

Dipartimento di Scienze Chimiche, Università di Napoli "Federico II", Italy
and Consorzio Interuniversitario di Reattività Chimica e Catalisi (CIRCC), Italy

Biodiesel, a mixture of fatty acid alkyl esters produced from vegetable oils (FAME), is a viable alternative to fossil fuels. It is conveniently obtained through simultaneous esterification and transesterification of vegetable oils with high acidity, such as frying oil residues or waste oil of other productions.¹



At industrial level, this process is typically promoted by alkaline catalysts, and one of the limits of this approach is their inevitable neutralization because of the acidity of the system. Therefore, the attention has been focused on catalysts Lewis acids, capable of promoting simultaneously the two reactions. However, it has been found that the use of these catalysts in homogeneous phase gives rise to problems related to their corrosive action, which makes the manufacture more complex. A solution to this drawback can derive from their heterogenization, which in addition to eliminate these problems, could favour separation and recycling. In this sense, an often winning strategy is the anchoring on solid supports of traditional homogeneous catalysts, according to the principles of so-called homogeneous supported catalysis.² In the light of these considerations, this study has selected as the most promising candidates complexes of Zn(II) which have nitrogen bidentate ligands with a functional group that allows to carry out the anchoring to a solid matrix (as SiO₂). The study of the supported complex **1** in the reaction of esterification and transesterification of a variety of oils, both neutral and with different contents of acidity, has revealed that the catalyst is active and can be recycled several times by simple decantation from the reaction mixture.




Figure 1.

References

- [1] Aresta, M.; Dibenedetto, A.; Dumeignil F. (Eds.), Biorefinery: from Biomass to Chemicals and Fuels, De Gruyter, **2012**.
[2] Li, C.; Liu, Y. (Eds) Bridging Heterogeneous and Homogeneous Catalysis: Concepts, Strategies, and Applications, Wiley-VCH, **2014**.

Synthesis and Cytotoxic Activity of Ruthenium Complexes with Ethacrynic Acid Derived Ligands

Gabriele Agonigi,^a Lorenzo Biancalana,^{**} Paul J. Dyson,^b Fabio Marchetti,^a Guido Pampaloni,^a Emilia Păunescu,^b Tina Riedel,^b Stefano Zacchini^c

^a Dipartimento di Chimica e Chimica Industriale, Università di Pisa, Via G. Moruzzi 13, 56124 Pisa (PI), Italy.


Email: lorenzo.biancalana@for.unipi.it

^b Institut des Sciences et Ingénierie Chimiques, Ecole Polytechnique Fédérale de Lausanne, Route Cantonale, 1015 Lausanne (VD), Switzerland.

^c Dipartimento di Chimica Industriale "Toso Montanari", Università di Bologna, Viale Risorgimento 4, 40136 Bologna (BO), Italy.

Decades of intensive research established the potential of some families of ruthenium compounds as anticancer agents. Thus, (LH)[trans-RuIIICl₄LL'] complexes, *i.e.* KP1019 (L = L' = κN -Indazole) and NAMI-A (L = κN -Imidazole, L' = κS -DMSO), are actually in later stages of clinical trials while several Ru(II)-arene complexes resulted to be very promising in pre-clinical trials.^{1,2} Metal complexes can be endowed with higher anticancer selectivity and effectiveness by incorporating, as ligands, organic fragments with complementary biological activity. Following a recent approach,³ we prepared new *P*- and *N*-donor ligands modified with Ethacrynic acid (EA-CO₂H). Ethacrynic acid, a diuretic in clinical use, is a potent inhibitor of Glutathione-S-Transferases (GSTs), a class of metabolic enzymes that are overexpressed in many tumors. GST enzymes are not only implicated in the development of drug resistance but also contribute to cancer progression via inhibition of programmed cell death.

In the present study, we synthesized novel EA-modified pyridine- and triphenylphosphine-based ligands, which were then used for the preparation of Ru(II)-*p*-cymene and Ru(III) NAMI-A type complexes. Ligands and complexes were fully characterized by analytical and spectroscopic methods and, in one case, by single-crystal X-ray diffraction. Moreover, some of the compounds underwent *in vitro* tests on ovarian carcinoma cell lines: the obtained results were similar to those previously reported for EA-functionalized Ru complexes.³



References

- [1] Weiss A.; Berndsen R. H.; Dubois, M.; Müller, C.; Schibli, R.; Griffioen, A. W.; Dyson P. J.; Nowak-Sliwinska P. *Chem. Sci.* **2014**, 5, 4742.
[2] Hartinger C. G.; Metzler-Nolte N.; Dyson P. J. *Organometallics* **2012**, 31, 5677.
[3] Ang W. H.; De Luca A.; Chapuis-Bernasconi C.; Juillerat-Jeanneret L.; Lo Bello M.; Dyson P. J. *ChemMedChem* **2007**, 2, 1799 and references therein.

Molecular Cages Based on Tris(2-pyridylmethyl)amine Complexes

Carlo Bravin,^{a*} Elena Badetti,^a Francesca A. Scaramuzzo,^a Rakesh Puttreddy,^b Kari Rissanen,^b Giulia Licini,^a Cristiano Zonta^a

^a Department of Chemical Sciences, University of Padova, via Marzolo 1, 35131 Padova.

Email: carlo.bravin@studenti.unipd.it

^b Department of Chemistry, University of Jyväskylä, P.O. Box 35, 40014 Jyväskylä, Finland

Biological and synthetic molecular containers have largely attracted the attention of scientists for their peculiar properties and applications.¹ In particular, supramolecular capsules and cages have offered in recent years the opportunity to draw important guidelines about the interpretation of the confinement phenomena. In this communication, we report the formation of a series of new self-assembled cages containing, in the inner part of the cavity, two metals with a coordination site available for binding. We have planned to take the advantage offered by dynamic covalent chemistry (DCC) for the synthesis of molecular cages of opportunely designed *tris*(2-pyridylmethyl)amine **TPy** metal complexes.² **TPy** complexes are known to furnish stable species that have been previously used for carboxylic acids recognition.³ The findings on their molecular recognition properties highlight unprecedented results in the thermodynamic of binding events that have a broad impact in the understanding of the molecular recognition phenomena.




Figure 1. Synthesis and X-ray crystal structure of the tris(2-pyridylmethyl)amine cage.

References

- [1] Ballester, P.; Fujita, M.; Rebek, J. *Chem. Soc. Rev.* **2015**, *44*, 392.
- [2] Scaramuzzo, F. A.; Licini, G.; Zonta, C. *Chem. Eur. J.* **2013**, *19*, 16809.
- [3] Joyce, L. A.; Maynor, M. S.; Dragna, J. M.; da Cruz, G. M.; Lynch, V. M.; Canary, J. W.; Anslyn, E.V. *J. Am. Chem. Soc.* **2011**, *133*, 13746.

Co-Based Heterogeneous Catalysts from Well-Defined Complexes: Synthesis and Applications

Dario Formenti,^{a*} Francesco Ferretti,^a Christoph Topf,^b Annette-Enrica Surkus,^b Kathrin Junge,^b Matthias Beller^b and Fabio Ragaini^a

^a Dipartimento di Chimica, Università degli Studi di Milano, Via C. Golgi 19, 20133 Milano (MI), Italy.

Email: dario.formenti@unimi.it

^b Leibniz-Institut für Katalyse e.V., Albert Einstein Strasse 29a, 18059 Rostock, Germany.

α -Diimines represent a family of nitrogen chelating ligands which found various applications in homogeneous catalysis.^{1,2} Here we report the first example of employing these ligands for the synthesis of Co-based supported heterogeneous catalysts starting from their well-defined complexes (Figure 1). The structure of these new materials, elucidated by XRD, XPS, TEM, Raman, TPD and TPR techniques, involves the simultaneous presence of metallic Co and Co oxides nanoparticles in contact with nitrogen-doped graphene (NDGr, derived by the pyrolysis of the nitrogen ligand) in a core-shell structure. These catalysts were successfully applied to the hydrogenation of aromatic nitro compounds. The reaction proceeds with a low catalyst loading (0.5 mol %, 10 mg/mmol) affording substituted anilines in high selectivities and yields. Furthermore, a close correlation with both the amount and the type of nitrogen in the NDGr-based catalytic material was established.³




Figure 1.

References


- [1] Viganò, M.; Ragaini, F.; Buonomenna, M. G.; Lariccia, R.; Caselli, A.; Gallo, E.; Cenini, S.; Jansen, J. C.; Drioli, E. *ChemCatChem* **2010**, *2*, 1150.
- [2] Cenini, S.; Ragaini, F.; Tollari, S.; Paone, D. *J. Am. Chem. Soc.* **1996**, *118*, 11964.
- [3] Formenti, D.; Ferretti, F.; Topf, C.; Junge, K.; Ragaini, F.; Beller, M., *manuscript in preparation*.

A Class of Novel Iron N-Heterocyclic Carbene Complexes for Redox Catalytic Reactions

Andrea Cingolani,^a Cristiana Cesari,^a Rita Mazzoni,^a Stefano Zacchini,^a Valerio Zanotti^a

^aDepartment of Industrial Chemistry "Toso Montanari", University of Bologna, Viale del Risorgimento 4, 40136 Bologna, Italy.
Email: andrea.cingolani6@unibo.it

Catalytic hydrogenation and transfer hydrogenation are among the most important reactions in organometallic catalysis. In recent years, cyclopentadienone iron complexes have drawn attention in this field of chemistry due to their air-water stability, availability from cheap starting materials, and unique catalytic features arising from the presence of a non-innocent ligand.¹ Herein, we report the synthesis and characterization of new carbonyl iron complexes combining different ligands, such as cyclopentadienones and N-heterocyclic carbenes (NHC). The unique features of both ligands have been exploited to obtain new metal-ligand bifunctional catalysts to be employed in hydrogenation and transfer hydrogenation, and to be compared with analogous ruthenium complexes previously investigated.²



Carbene coordination occurred through a straightforward transmetallation reaction from the silver carbene intermediate (prepared *in situ* from the corresponding imidazolium salts) to the dicarbonyl acetonitrile iron intermediate (obtained *in situ* from the corresponding tricarbonyl complex) leading to complexes **1** and **2**.

Moreover, a microwaved-assisted method for the synthesis of tricarbonyl precursors has been developed. Microwave conditions resulted in shorter reaction time respect to the reflux reaction, without decreasing the yield.

All the complexes have been characterized by spectroscopy (NMR, IR, ESI-MS) and X-ray studies.

References

- [1] A. Quintard and J. Rodriguez *Angew. Chem. Int. Ed.* **53** (2014), 4044.
 [2] C. Cesari, S. Conti, S. Zacchini, V. Zanotti, M.C. Cassani and R. Mazzoni *Dalton Trans.* **43** (2014), 17240

Ruthenium(0) Cyclopentadienone N-Heterocyclic Carbene Complexes as Transfer Hydrogenation Catalyst: Synthesis and Reactivity

Cristiana Cesari,^{*,a} Andrea Cingolani,^a Rita Mazzoni,^a Chiara Parise,^a Stefano Zacchini,^a Valerio Zanotti^a

^a Department of Industrial Chemistry "Toso Montanari", Viale del Risorgimento 4, 40136 Bologna (BO), Italy.
Email: cristiana.cesari2@unibo.it

N-heterocyclic carbenes (NHC) are efficient ancillary ligands due to their strong coordination ability and their tuneable character which allows control of the steric and electronic properties of the metal centre.¹ Herein we report the synthesis, characterization and catalytic application of new carbonyl ruthenium complexes that containing tetraphenylcyclopentadienone and NHC ligands.²

The catalytic activity of the imidazolylienes (**1**) and triazolylidenes (**2**) ruthenium complexes was investigated in homogeneous catalysis: the complexes **1a** ($R' = R'' = CH_3$), **1b** ($R' = Bu, R'' = py$) and **2a** were tested as catalyst precursors for hydrogen transfer reactions (Figure 1).




Figure 1.

Transfer hydrogenation of ketones is catalyzed by the ruthenium(0) carbene complexes of type **1** and **2** upon treatment with cerium(IV) ammonium nitrate (CAN) as oxidant.³ Since the oxidation of the metal center seems to be the key issue for the catalyst activation, a formal oxidation to ruthenium(II) was also obtained by protonation of **1a** and **1b** with trifluoromethanesulfonic acid.

References

- [1] Hopkinson, M.N.; Richter, C.; Schedler, M.; Glorius, F.; *Nature*, **2014**, *510*, 485-496.
 [2] Cesari, C.; Conti, S.; Zacchini, S.; Zanotti, V.; Cassani, M. C.; Mazzoni, R.; *Dalton Trans.*, **2014**, *43*, 17240-17243.
 [3] Cesari, C.; Mazzoni, R.; Muller-Bunz, H.; Albrecht, M.; *J. Organomet. Chem.* **2015** <http://dx.doi.org/10.1016/j.jorganchem.2015.02.022>.


Poster 69

Catalytic CO₂ Hydroboration with Novel CO₂ Adducts of N/Si+ Frustrated Lewis Pairs

Niklas von Wolff,^a Guillaume Lefèvre,^a Jean-Claude Berthet,^a Pierre Thuéry,^a Thibault Cantat^{a*}


^a Laboratoire de Chimie Moléculaire et de Catalyse pour l'Energie, CEA Saclay, UMR 3685 CEA/CNRS, 91191, Gif-sur-Yvette.
Email: thibault.cantat@cea.fr, niklas.von-wolff@cea.fr

In recent years, cationic +IV silicon species (silyliums) have found application in the design of frustrated Lewis pairs (FLPs) for the activation of small molecules.¹ Highly electrophilic cationic silylium species however are strong σ and π acceptors and tend to undergo unwanted side-reactions (e.g. with solvents). Therefore the use of base-stabilized silyl cations with tunable reactivity is of interest. Although CO₂ inserts readily into N-Si bonds,² up today the reactivity of nitrogen/silylium FLPs for the reductive transformation of CO₂ and the activation of other small molecules has not yet been explored.



Scheme 1.

Using a silyl cation stabilized by an intramolecular nitrogen base (Scheme 1), CO₂ can be added to form an isolable N/Si+ FLP-CO₂ adduct. This system was further shown to catalyze the hydroboration of CO₂³ to methoxyboranes. The influence of the nature of the Si-R groups, as well as the counter-ion on the CO₂-adduct stability and the catalytic performance was investigated and suggest different mechanisms for the catalytic hydroboration of CO₂ depending on the hydride source (Scheme 2). DFT calculations and the isolation of a proposed key intermediate adumbrate a novel electrophilic activation of pinacolborane to form highly reactive base-stabilized silanes and highlight the role of CO₂ adducts during the catalytic cycle of CO₂ hydroboration.



Scheme 2.

References

- [1] T. Müller et al., *Angew. Chem. Int. Ed.*, **2011**, 12636; T. Müller et al., *Organometallics*, **2013**, 6736; A. E. Ashley et al., *Chem. Comm.* **2014**, 12753; T. Müller et al. *Chem. Eur. J.* **2014**, 9381.
- [2] E. A. V. Ebsworth, G. Rocktäschel, J. C. Thompson, *J. Chem. Soc. (A)* **1967**, 362; L. Birkhofer et al. *J. Organomet. Chem.* **1975**, 99, C1-C4; W. Kaim et al. *J. Organomet. Chem.* **1995**, 283
- [3] T. Cantat et al., *Chem. Eur. J.* **2014**, 10708; M.J. Sgro, D.W. Stephan, *Angew. Chem. Int. Ed.*, **2012**, 51, 11343-11345; F. G. Fontaine et al., *J. Am. Chem. Soc.*, **2013**, 9326; F. G. Fontaine et al., *J. Am. Chem. Soc.*, **2014**, 10708; D. W. Stephan, G. Ménard, *J. Am. Chem. Soc.*, **2010**, 1796; S. Sabo-Etienne et al., *Angew. Chem. Int. Ed.* **2012**.

Poster 70

Stereoselective Cyclopropanation of Olefins Promoted by a C₂-Symmetrical Iron(III) Porphyrin Complexes.


Daniela M. Carminati,^{a*} Daniela Intrieri,^a Stéphane Le Gac,^b Bernard Boitrel,^b Emma Gallo^a

^aUniversità degli Studi di Milano, via C. Golgi 19, 20133 Milano, Italy.

Email: daniela.carminati@unimi.it

^bInstitut des Sciences Chimiques de Rennes, Université de Rennes 1, 263 avenue du Général Leclerc, 35042 Rennes Cedex, France.

Cyclopropanes are important three-member carbon rings which often represent the active part in biological and/or pharmaceutical compounds.¹⁻² Catalytic diastereo- and enantioselective olefin cyclopropanations have been extensively explored and among all the available synthetic methodologies to synthesise cyclopropanes, the one pot reaction of diazocompounds with olefins represents a sustainable and atom-efficient strategy due to the formation of N₂ as the only stoichiometric by-product. Considering that metal porphyrins represent a very competent and stereoselective class of catalysts, we are currently studying the catalytic activity of iron derivatives which display a good sustainability for the presence of the eco-friendly, cheap and very efficient metal centre. Thus, we synthesised the new iron(III) complex **1Fe** starting from the porphyrin ligand **1**³ (scheme 1) which has one C₂ axis within the porphyrin plane and exhibits an open space on each side for the substrate access. Complex **1Fe** demonstrated a very high catalytic activity in cyclopropanations which occurred with excellent *trans*-diastereoselectivities (94-99%). Since positive data, we employed ligand **1** to synthesise the chiral derivative **2**, which shows a steric chiral bulk surrounding the *N*-core of the porphyrin. The free porphyrin **2** was then reacted with iron bromide, in the presence of methanol, to yield the relative iron(III) complex **2Fe** (scheme 1) which was completely characterised.⁴ Complex **2Fe** showed a very high catalytic efficiency; excellent *trans*-diastereoselectivities, good enantioselectivities and outstanding TON and TOF values were observed in the reaction of cyclopropanation of several olefins by differently substituted diazo compounds (scheme 1).⁴



Scheme 1.

References

- [1] C. A. Carson, M. A. Kerr, *Chem. Soc. Rev.* **2009**, 38, 3051-3060.
- [2] A. K. Kumar, *Int. J. Pharm. Pharm. Sci.* **2013**, 5, 467-472.
- [3] A. Didier, L. Michaudet, D. Ricard, V. Baveux-Chambenoît, P. Richard, B. Boitrel, *Eur. J. Org. Chem.* **2001**, 1927-1926.
- [4] D. Intrieri, S. Le Gac, A. Caselli, E. Rose, B. Boitrel, E. Gallo, *Chem. Commun.* **2014**, 50, 1811-1813.

C-H Activation Reactions with Rhodium(III) Complexes Containing Tridentate Ligands

Filippo Brusa,* Umberto Piarulli, Gianluigi Brogгинi

Università dell'Insubria, Dipartimento di Scienza e Alta Tecnologia, Via Valleggio 11, 22100 Como, Italy, *filippo.brusa@uninsubria.it

Over the last few years, various ligands containing chiral dihydrooxazole rings as coordinating units have been reported, and many of them have found successful applications in asymmetric catalysis.¹ A common feature of these ligands is their rather rigid framework with two stereogenic centers in close proximity to the coordination sphere. These ligands are quite attractive since the dihydrooxazole units are easily prepared from enantiomerically pure amino-alcohols, that are commercially available or can be readily prepared from amino acids. More recently, the synthesis of a chiral tridentate ligand containing two dihydrooxazole rings has been reported² and its coordination to Zn^{II}, Ni^{II} and Cu^{II} studied.

As part of our ongoing interest in transition metal mediated heterocyclization reactions,³ we have prepared a new rhodium(III) complex (Figure 1) bearing two chiral dihydrooxazole rings, whose reactivity and selectivity has been studied in different benchmark reactions.




Figure 1.

References

- [1] Pfaltz A. *Acta Chem. Scand.* **1996**, *50*, 189.
- [2] Fahrni C. J.; Pfaltz A. *Helv. Chim. Acta* **1998**, *81*, 491.
- [3] Borsini, E.; Broggin, G.; Fasana, A.; Galli, S.; Khansaa, M.; Piarulli, U.; Rigamonti, M. *Adv. Synth. Catal.* **2011**, *353*, 985.

Glycoporphyrin Catalysts for Efficient C-H Bond Aminations by Organic Azides

Giorgio Tseberlidis,* Paolo Zardi, Alessandro Caselli and Emma Gallo

^a Department of Chemistry, University of Milan, Via Golgi 19, 20133 Milan, Italy. Email: giorgio.tseberlidis@unimi.it

The activation of C-H bonds represents a 'hot' topic due to the enormous synthetic potential of this class of organic transformations and the ubiquity of C-H functionalities in almost every molecular scaffold. Among all kinds of C-H bond activations, the C-H bond amination allows the conversion of low cost products, such as simple hydrocarbons, into high added value nitrogen-containing derivatives which often present important biological and/or pharmaceutical features.¹ Amongst all the catalysts active in C-H bond aminations, metal porphyrins have a good catalytic efficiency as well as an acceptable biocompatibility.²⁻⁵ In order to enhance this latter catalyst characteristic it is important to introduce favourable functional groups (such as carbohydrates) on the porphyrin skeleton and low toxicity transition metals into the porphyrin core. We report herein the synthesis of new glycoporphyrin ligands which bear a glucopyranoside derivative on each *meso*-aryl moiety of the porphyrin skeleton. The saccharide unit is either directly conjugated to the porphyrin or a triazole spacer is placed in between the carbohydrate and porphyrin ring. The obtained glycoporphyrin ligands were employed to synthesise cobalt(II), ruthenium(II) and iron(III) complexes which were tested as biocompatible catalysts of C-H bond aminations by organic azides. Two of the synthesised complexes were very efficient in promoting catalytic reactions, up to 19 compounds were isolated in good to excellent yields, and achieved results indicated that ruthenium and iron complexes show an interesting complementary catalytic activity in several amination reactions. The eco-friendly iron catalyst displayed very good chemical stability catalysing the amination reaction for three consecutive runs without losing the catalytic activity.




Figure 1.

References

- [1] Yamaguchi, J.; Yamaguchi, A. D.; Itami, K. *Angew. Chem. Int. Ed. Engl.* **2012**, *51*, 8960-9009.
- [2] Lu, H.; Zhang, X. P. *Chem. Soc. Rev.* **2011**, *40*, 1899-1909.
- [3] Che, C.-M.; Lo, V. K.-Y.; Zhou, C.-Y.; Huang, J.-S. *Chem. Soc. Rev.* **2011**, *40*, 1950-1975.
- [4] Fantauzzi, S.; Caselli, A.; Gallo, E. *Dalton Trans.* **2009**, 5434-5443.
- [5] Oszajca, M.; Franke, A.; Brindell, M.; Stochel, G.; van Eldik, R. *Coord. Chem. Rev.* **2015**, Ahead of Print.

Poster 73

Diffusion-Ordered Spectroscopy (DOSY) : a Simple, Promising Tool for Analysing Complex Reaction Systems

Ricky Payen,^{a*} Cécile Le Duff,^a Olivier Riant,^a Tom Leyssens^a

^a Institute of Condensed Matter and Nanosciences, Université catholique de Louvain, 1348 Louvain-la-Neuve, Belgium.

Email: ricky.payen@uclouvain.be

Diffusion-Ordered Spectroscopy (DOSY) is also known as the Nuclear Magnetic Resonance (NMR) chromatography since it allows the separation of a mixture of compounds spectroscopically. It is a simple technique which only requires classical NMR hardware and software. In this work, we report the use of DOSY as a promising tool to estimate diffusion coefficients of copper (I) complexes in solution.

Copper (I) dibenzoylmethane (DBM) complexes are characterized as monomeric species in the solid state. Binap- and Segphos- copper (I) chloride complexes are reported as dimers in the solid state. Their nature in solution is uncertain.¹




Figure 1. Single crystal XRD structures of dpephosCuDBM and [BinapCuCl]₂ complexes

Using the Stokes-Einstein modified equation,² it is possible to estimate the diffusion coefficients theoretically. Comparing these values to experimental data, we confirmed the monomeric nature of diphosphine copper (I) DBM complexes and dimeric nature of diphosphine copper (I) chloride complexes in toluene. This method can be applied in the investigation of reaction mechanisms involving these copper (I) complexes.

References

- [1] G. Hattori, K. Sakata, H. Matsuzawa, Y. Tanabe, Y. Miyake, Y. Nishibayashi, *Journal of the American Chemical Society* **2010**, *132*, 10592-10608
- [2] R. Evans, Z. Deng, A. K. Rogerson, A. S. McLachlan, J. J. Richards, M. Nilsson, G. A. Morris, *Angewandte Chemie International Edition* **2013**, *52*, 3199-3202.

Poster 74

Synthesis and Reactivity of Ni-P Carbonyl Clusters

Chiara Capacci,^{a*} Maria Carmela Iapalucci,^a Cristina Femoni^a

^a School of Science, Department of Industrial Chemistry "Toso Montanari", University of Bologna, Viale del Risorgimento 4, 40136 Bologna (BO), Italy, *chiara.capacci5@unibo.it

Over the past years Ni-P supported nanoparticles have been used as catalysts for many reactions,¹⁻² and it is known that carbonyl clusters can be used as precursor for the synthesis metallic nanoparticles.³

Therefore, we chose to synthesize new carbonyl species containing nickel and phosphorus. There were no homoleptic carbonyl clusters known in the literature and only examples with phosphines as ligands were reported.⁴

The reaction between the [Ni₆(CO)₁₂]²⁻ carbonyl cluster and PCl₃ or POCl₃ in different conditions allowed to obtain new species, such as [Ni₁₁P(CO)₁₈]³⁻, [Ni₁₄P₂(CO)₂₂]²⁻, [Ni_{23-x}P₂(CO)_{30-x}]⁴⁻ (x=0, 1), [HNi₃₁P₄(CO)₃₉]⁵⁻. Once available, we studied their reactivity and the results showed that these species are strictly related: under the appropriate conditions they can convert one into another (Figure 1).

Finally, we tried to investigate the catalytic properties of these Ni-P compounds, but the studies we carried out have not shown relevant results yet. However it has been confirmed (via TEM analyses) that the controlled thermal decomposition of [HNi₃₁P₄(CO)₃₉]⁵⁻ leads to the formation of heteroatomical nanoparticles.




Figure 1.

References

- [1] Sh. F. Yang, Ch. H. Liang, R. A. Prins; *J. Catal.*, **2006**, *237*, 118.
- [2] S. J. Sawhill, K. A. Layman, D. R. Van Wyk, M. H. Engelhard, Ch. M. Wang; *J. Catal.*, **2005**, *231*, 300.
- [3] S. Albonetti, C. Femoni, R. Bonelli, J. E. Mengou, C. Tiozzo, S. Zacchini, F. Trifirò; *Catalysis today*, **2008**, *137*, 483-488.
- [4] P. D. Mlynek, L. F. Dahl; *Organometallics*, **1997**, *16*, 1655.

¹⁵N NMR Spectroscopy Unambiguously Establishes the Coordination Mode of the Diimine Linker 2-(2'-Pyridyl)Pyrimidine-4-Carboxylic Acid (CpPh) in Ru(II) Complexes

Federica Battistin,^{a*} Gabriele Balducci,^a Nicola Demitri,^b Elisabetta Inengo,^a Barbara Milani,^a Enzo Alessio^a

^a Department of Chemical and Pharmaceutical Sciences, Università di Trieste, Via L. Giorgieri 1, 34127 Trieste, Italy.

Email: federica.battistin@phd.units.it

^b Elettra – Sincrotrone Trieste, S.S. 14 Km 163.5 in Area Science Park, 34149 Basovizza – Trieste, Italy.

In 2009 Spiccia and coworkers introduced a new diimine linker, 2-(2'-pyridyl)pyrimidine-4-carboxylic acid (cpPh, Figure 1), as an easy-to-make alternative to the widely used – but more tedious to prepare – 4'-methyl-2,2'-bipyridine-4-carboxylic acid (bpyAc, Figure 1).²




Figure 1. The diimine ligands bpyAc and cpPh with numbering schemes and labels.

This linker can originate linkage isomers, in fact its pyrimidine ring can bind to the metal ion either through the nitrogen atom *ortho* (N^o) or *para* (N^p) to the carboxylate linked to C4. Most of the cpPh-Ru(II) conjugates reported so far were prepared following a synthetic route that led selectively to the N^p coordination mode.^{1,3-6}

In this work we investigated the reactivity of the Ru(II) precursor *trans,cis,cis*-[RuCl₂(CO)₂(dmsO)₂] towards cpPh. We isolated and characterized two isomers and we reported how ¹H, ¹⁵N- HMBC NMR spectra recorded at natural abundance of ¹⁵N isotope on these compounds allowed us to unambiguously establish the binding mode of cpPh, also in the absence of the X-ray structure.

References

- [1] Nickita, N.; Gasser, G.; Pearson, P.; Belousoff, M. J.; Goh, L. Y.; Bond, A. M.; Deacon, G. B.; Spiccia, L. *Inorg. Chem.* **2009**, *48*, 68-81.
- [2] McCafferty, D. G.; Bishop, B. M.; Wall, C. G.; Hughes, S. G.; Mecklenberg, S. L.; Meyer T. J.; Erickson, B. W. *Tetrahedron* **1995**, *51*, 1093-1106.
- [3] (a) Joshi, T.; Barbante, G. J.; Francis, P. S.; Hogan, C. F.; Bond, A. M.; Gasser, G.; Spiccia, L. *Inorg. Chem.* **2012**, *51*, 3302-3315; (b) Joshi, T.; Gasser, G.; Martin, L. L.; Spiccia, L. *RSC Adv.* **2012**, *2*, 4703-4712; (c) Joshi, T.; Patra, M.; Spiccia, L.; Gasser, G. *Artificial DNA: PNA & XNA* **2013**, *4*, 11-18. (d) Bischof, C.; Joshi, T.; Dimri, A.; Spiccia, L.; Schatzschneider, U. *Inorg. Chem.* **2013**, *52*, 9297-9308.
- [4] Joshi, T.; Pierroz, V.; Ferrari, S.; Gasser, G. *ChemMedChem* **2014**, *9*, 1231-1237.
- [5] Pierroz, V.; Joshi, T.; Leonidova, A.; Mari, C.; Schur, J.; Ott, I.; Spiccia, L.; Ferrari, S.; Gasser, G. *J. Am. Chem. Soc.* **2012**, *134*, 20376-20387.
- [6] Joshi, T.; Pierroz, V.; Mari, C.; Gemperle, L.; Ferrari, S.; Gasser, G. *Angew. Chem. Int. Ed.* **2014**, *53*, 2960-2963.

Copper(II) Complexes of Arylhydrazone of 1*H*-Indene-1,3(2*H*)-Dione as Catalysts for the Oxidation of Alcohols


Gonçalo A. O. Tiago,^{a,*} Ana P. C. Ribeiro,^a Luis Branco,^b Kamran T. Mahmudov,^{a,c}
M. Fátima C. Guedes da Silva,^a Armando J. L. Pombeiro^a

^a Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001, Lisbon, Portugal.

E-mail: goncalo.tiago@ist.utl.pt

^b REQUIMTE, Departamento de Química, Faculdade de Ciências e Tecnologia da Universidade Nova de Lisboa, Quinta da Torre, Caparica, Portugal. ^c Department of Chemistry, Baku State University, Z. Xalilov Str. 23, Az 1148 Baku, Azerbaijan.

Copper(II) acetate was allowed to react with arylhydrazone of 1*H*-indene-1,3(2*H*)-dione – sodium 2-(2-(1,3-dioxo-1*H*-inden-2(3*H*)-ylidene)hydrazinyl)benzenesulfonate (**NaHL**; see Figure) in the absence and in the presence of diethanolamine, leading to the formation of complexes **1** and **2** which were characterized by conventional methods, as well as by single crystal X-ray diffraction.



Both complexes act as active and selective catalysts for the solvent free oxidation of 1-phenylethanol to acetophenone, using *tert*-butylhydroperoxide (TBHP) as oxidant, under mild conditions. Complex **2** exhibits a higher activity than **1**. An alternative medium [1] (ionic liquid) was also tested and the differences in yield and reaction time are discussed.

Acknowledgements: This work has been partially supported by the Foundation for Science and Technology (FCT), Portugal (UID/QUI/00100/2013). G.A.O.T. is also thankful to the CATSUS doctoral FCT program for his PhD fellowship. The authors acknowledge the Portuguese NMR Network (IST-UL Centre) for access to the NMR facility and the IST Node of the Portuguese Network of Mass-spectrometry for the ESI-MS measurements.

References

- [1] R. Jlassi, A.P.C. Ribeiro, M.F.C.G. da Silva, K. T. Mahmudov, M. N. Kopylovich, T.B. Anisimova, H. Naili, G.A.O. Tiago, and A.J.L. Pombeiro, "Polynuclear Copper(II) Complexes as Catalysts for the Peroxidative Oxidation of Cyclohexane in a Room-Temperature Ionic Liquid", *Eur. J. Inorg. Chem.*, DOI:10.1002/ejic.201402352.

Poster 77


Divergent Intramolecular Hydroamination Reactions of Terminal Alkynes Under Palladium and Platinum Catalysis

Alberto Mazza, Egle M. Beccalli

DISFARM, Sez. Chimica Generale e Organica "A. Marchesini", Università degli Studi di Milano, Via Venezian 21, 20133 Milano.
Email: alberto.mazza@unimi.it

Transition-metal-catalyzed intramolecular reactions involving a C-C multiple bonds have been appreciated as useful processes to obtain heterocyclic systems.¹ Among them palladium in particular has been played a prominent role. For a long time our research group was studying the palladium-catalyzed intramolecular reactions and functionalizations to obtain polycyclic systems² and in this contest the hydroamination reactions involving the C-N bonds formation continue to gain our interests.

The present work is regarding the reactivity of terminal alkynes and alkenes tethered to a nucleophilic pendant as a useful scaffolds for intramolecular hydroamination processes with the aim to obtain aza-containing systems. The use of different transition metal-catalysts results in a complete divergent *endo/exo* selectivity depending on the metal catalyst used: in the presence of Pd(PPh₃)₄ 6-*exo*-dig cyclization was observed, while with PtCl₂ the 7-*endo*-dig cyclization occurred. Moreover working on the allyl derivatives under the same platinum catalysis, the dihydrobenzoxazine rings formation was reported.



References

- [1] a) Nakamura, I.; Yamamoto, Y., *Chem. Rev.* **2004**, *104*, 2127. b) Zhang, M., *Adv. Synth. Catal.*, **2009**, *351*, 2243. c) Magano, J.; Dunetz, J. R., *Chem. Rev.*, **2011**, *111*, 2177.
[2] a) Beccalli, E. M. et al. *Adv. Synth. Cat.* **2013**, *355*, 1640. b) Beccalli, E. M. et al. *J. Organomet. Chem.* **2010**, *696*, 277. c) Beccalli, E. M. et al. *J. Org. Chem.* **2010**, *75*, 6923. d) Beccalli, E. M. et al. *Tetrahedron* **2009**, *65*, 3486. e) Beccalli, E. M. et al. *J. Org. Chem.* **2008**, *73*, 4746.

Poster 78


The bimca-ligand: Synthesis and Application of its Rhodium and Iridium Complexes

Eva Jürgens,^a Alexander Seyboldt,^a Doris Kunz^a

^aInstitute for Inorganic Chemistry, University of Tübingen, Auf der Morgenstelle 18, 72074 Tübingen
Email: eva.juergens@uni-tuebingen.de

Introduced by Shaw in 1976, pincer-type ligands have become an important ligand class in organometallic chemistry during the last decades.¹ Nearly 15 years ago, they were combined with NHCs either as backbone or side donor moieties. Nowadays a broad range of tridentate mono- or biscarbene ligands are increasingly used in the synthesis of highly active homogenous catalysts.²

In 2007 we reported the monoanionic bis(imidazolynilidene)carbazolide (bimca) ligand as anionic pincerligand with two carbene moieties.³ Since then we established a simplified synthetic access to its precursor and prepared a decent-sized ligand library.⁴ Reaction of the lithium carbenes with Rh(I) carbonyl precursors led to the formation of a Rh (bimca)(CO) complex **1a**.



This complex shows a highly nucleophilic reactivity caused by the strong σ -donor and weak π -acceptor character of the NHC moieties in addition to the monoanionic carbazolide donor.³⁻⁵ **1a** could be applied in the catalytic rearrangement of epoxides to methyl ketones (*Meinwald* rearrangement), using LiNTf₂ as cocatalyst for a preactivation of the epoxides.⁵

References

- [1] Pugh, D.; Danopoulos, A. *Coord. Chem. Rev.* **2007**, *251*, 610–641. [1b] Albrecht, M.; van Koten, G.; *Angew. Chem. Int. Ed.* **2001**, *40*, 3750–3781.
[2] Jürgens, E.; Kunz, D. in *Molecular Catalysts* **2014**, 183–206.
[3] Moser, M.; Wucher, B.; Rominger, F.; Kunz, D. *Organometallics* **2007**, *26*, 1024–1030.
[4] *Manuscript in*
[5] Jürgens, E.; Wucher, B.; Rominger, F.; Törnroos, K. W.; Kunz, D. *Chem. Commun.* **2015**, *51*, 1897–1900.

Poster 79

Synthesis of Metal Nanoparticles by Bio-Based Derivatives

Andrea Mascitti,^{a*} Nicola d'Alessandro,^a Lucia Tonucci^b

^a Department of Engineering and Geology (INGEO), "G.d'Annunzio" University of Chieti-Pescara, Viale Pindaro, 42, 65127 Pescara, Italy.

Email: andrea.mascitti@studenti.unich.it

^b Department of Philosophical, Educational and Economic Sciences, "G.d'Annunzio" University of Chieti-Pescara, Via dei Vestini 31, 66100 Chieti, Italy.

Benzoxaboroles In view of new and important commercial opportunities, it is important to meet the green chemistry principles for the overall procedure of the preparation of metal NPs, minimizing reactant consumption and by-product formation, and using, if possible, renewable materials and benign solvents.

Indeed, in the past, some studies have reported the formation of Au, Ag, Pt and Pd NPs by natural products, like starch, carbohydrates, wood nanomaterials, cellulose nanocrystallites, hydrocolloid gums and various plant extracts¹. In other studies, NPs were synthesized by the polyol process, where a polyalcohol, generally coupled with a base, acts not only as a solvent but also as a mild reducing agent for the metal salt precursors². In this context, glycerol has successfully employed in the preparation of a variety of metal NPs. However, although the preparation of transition metals nanoparticles has been extensively studied, the literature contains relatively few references on the synthesis of nanoparticles of group 7 transition metals, namely Mn, Tc and Re.

Our main interest in this study was addressed to the green synthesis of rhenium NPs. One of the main drawbacks in the development of a method for the synthesis and also for the potential applications, of Re NPs, is the low stability of the Re colloids generated by chemical reductions.

Our synthetic strategy was based on the reduction of ReCl₅ to Re NPs by use of lignin as reductant and stabilizing agent and glycerol or water as media of the reaction; we have to point out that lignin and glycerol are two of the most widely diffused by-products of the agro-industry. For a comparison purpose, we initially synthesized Re NPs employing NaBH₄ as reductant and rhenium perrhenate as starting Re source: the reaction was conducted in water or water/glycerol solution. The green synthetic procedures were then developed starting from a water or glycerol solution containing lignin as reductant and rhenium (V) chloride as metal source. In a first protocol, such solution was heated at 80 °C for 3 h while in a second protocol the reaction mixture was sonicated (20 kHz) and heated at 60 °C for 3 h. All the Re NPs were characterized using TEM, UV-vis, XRD analysis. During the characterization of NPs, we noted that glycerol is not only a solvent but it contributes to stabilize the nanoparticles in a synergic effect with lignin.

At the moment, we are studying the application of these nanoparticles, for example in the reduction of succinic acid to 1,4-butadienol.

References

- [1] Raveendran, P.; Fu J.; Wallen, S. L. *J. Am. Chem. Soc.* **2003**, *125*, 13940.
- [2] Carroll, K.J.; Reveles, J.U.; Shultz, M.D.; Khanna, S.N.; Carpenter, E.E. *J. Phys. Chem., C* **2011**, *115*, 2656.
- [3] Coccia, F.; Tonucci, L.; Bosco, D.; Bressan, M.; d'Alessandro, N. *Green Chem.* **2012**, *14*, 1073.

Poster 80

Synthesis, Biological Evaluation and Stability Studies of Novel Symmetrical and Non-Symmetrical 11th Group-NHC Complexes

Marika Marinelli,^a Maura Pellei,^a Cristina Marzano,^b Valentina Gandin,^b Alessandro Dolmella,^b H. V. Rasika Dias^c and Carlo Santini^a

^a School of Science and Technology, Chemistry Division, University of Camerino, via S. Agostino 1, 62032 Camerino, Italy.

Email: marika.marinelli@unicam.it

^b Department of Pharmaceutical and Pharmacological Sciences, University of Padova, via Marzolo 5, Padova, Italy.

^c Department of Chemistry and Biochemistry, The University of Texas at Arlington, Box 19065, Arlington, Texas 76019-0065, United States.

Recently metallodrug chemistry bearing N-heterocyclic carbene (NHC) supporting ligands has experienced a new rebirth. Our research group, during the last 7 years, has developed different classes of coinage metal-NHC complexes obtained from the precursors {[H₂B(BnTzH)₂]Br},¹ {[H₂B(pNO₂BnTzH)₂]Br} {[HTz(pNO₂Bn)₂]Br}². Recently we have focused the research work on the synthesis of new coinage metal-NHCs complexes obtained from the water soluble ligands Him^{1R,3R}Cl (R = COOCH₃, COOCH₂CH₃ or CON(CH₂CH₃)₂)³ or the zwitterionic symmetrical NHC ligands NaHim^{1R,3R,4R} (R = (CH₂)₃SO₃⁻, R' = H, CH₃)⁴ (Figure 1), NaHBzim^{1R,3R} (R = (CH₂)₃SO₃⁻) and non-symmetrical NHC ligands NaHim^{1R,3R} (R = (CH₂)₃SO₃⁻, R' = CH₂C₆H₅), {[HBzim^{1R,3R}]Br} (R = (CH₂)₃SO₃⁻Na, R' = CH₂C₆H₅) and the related copper(I)- and silver(I)-NHC complexes.⁵ Finally we studied the cytotoxic activity and the stability of the NHC complex classes developed by our research group.




Figure 1.

References

- [1] Papini, G.; Bandoli, G.; Dolmella, A.; Gioia Lobbia, G.; Pellei, M.; Santini, C., *Inorg. Chem. Commun.* **2008**, *11*, 1103.
- [2] Marinelli, M. *et al.*, unpublished results.
- [3] Papini, G.; Pellei, M.; Gioia Lobbia G.; Burini, A.; Santini, C., *Dalton Trans.* **2009**, 6985.
- [4] Pellei, M.; Gandin, V.; Marinelli, M.; Marzano, C.; Yousufuddin, M.; Dias, H. V. R.; Santini, C., *Inorg. Chem.* **2012**, *51*, 9873.
- [5] Gandin, V.; Pellei, M.; Marinelli, M.; Marzano, C.; Dolmella, A.; Giorgetti, M.; and Santini, C., *J. Inorg. Biochem.* **2013**, *129*, 135-

Poly(Alkylideneamine)s Dendrimers with Nitrile and Amine Terminal Groups as a Platform to Deliver Metallo drugs

Dina Maciel,^a María de los Ángeles Muñoz-Fernández,^b Helena Tomás,^a João Rodrigues^{a*}

^a CQM – Centro de Química da Madeira, MMRG, Universidade da Madeira, Campus Universitário da Penteada, 9020-105 Funchal, Portugal.

Email: joaor@uma.pt

^b Laboratorio de Inmunobiología Molecular, Hospital General Universitario Gregorio Marañón, Madrid, Spain

The acquired immunodeficiency syndrome (AIDS) infected around 78 million of people since the beginning of the epidemic.¹ On the other hand, cancer is one of the leading causes of deaths worldwide. Metal complexes, like ruthenium compounds, present less toxicity when comparing to platinum derivatives and can bind to proteins in the plasma due to their ability to mimic the behavior of iron.² The combination of metals with dendrimers, which are highly branched molecules with well-defined structures, provides new nanostructures (metallo dendrimers) with improved biological properties.^{3,4} These metallo dendrimers are under study to be used in a broad range of applications including in the biomedical field to fight cancer and infections.

The aim of this project is to develop a new family of metallo dendrimers based on poly(alkylideneamine) dendrimers⁵ and ruthenium or platinum complexes, with improved solubility in biological media, and to evaluate their potential as effective drugs against HIV-1 infection/replication and different types of cancer.

In this work, the preliminary results on the preparation and characterization, by NMR, IR and MS, of this new family of poly(alkylideneamine)-based dendrimers having, at each generation, nitrile and amine terminal groups, will be presented and discussed. The nitrile and amine terminal groups on the surface of dendrimers will serve to grow the dendrimer generation and will act as bridging groups for complexation of metallo drugs.

Acknowledgments: This research was supported by Fundação para a Ciência e a Tecnologia (FCT) with Portuguese Government funds through the CQM Strategic Project PEst-OE/QUI/UI0674/2014, the NMR Network – (PTNMR-2014 and PTNMR-2015) and the PhD Grant SFRH/BD/102123/2014 (D.M). The support of the international network CYTED 214RT0482 in the domain of the HIV infection is highly appreciated. We also acknowledge the continuous support of our work by Hotel Vidamar Resorts Madeira.

References

- [1] <http://www.who.int/gho/hiv/en/> - accessed on 29 of July.
- [2] Smith, G.S.; Therrien, B. *Dalton Trans.* **2011**, *40*, 10793.
- [3] Govender, P.; Renfrew, A.K.; Clavel, C.M.; Dyson, P.J.; Therrien, B.; Smith, G.S. *Dalton Trans.* **2011**, *40*, 1158.
- [4] Viñas, C.; Teixidor, F.; Núñez, R. *Inorg. Chim. Acta.* **2014**, *409*, 12.
- [5] a) Jardim, M.G.; Rissanen, K.; Rodrigues, J. *Eur. J. Inorg. Chem.* **2010**, 1729. b) Rodrigues J.; Jardim M. J.; Gouveia M.; Tomás H.; Rissanen K., *New J. Chem.* **2011**, *35*, 1938.

Metal Derivatives with 3-(2-sulfonic phenylhydrazo)pentane-2,4-dione, and 3,7-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane: Synthesis and Characterization

A.G. Mahmoud,^{a,*} K.T. Mahmudov,^a M.F.C. Guedes da Silva,^a M.J.F. Calvete,^b A.J.L. Pombeiro^a

^a Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001, Lisbon, Portugal.

E-mail: abdallah1126@gmail.com

^b Department of Chemistry, University of Coimbra, Rua Larga, 3004-535 Coimbra, Portugal

As part of our interest on new water soluble complexes, new set of Transition metal complexes were prepared using two water soluble ligands, which are Sodium salt of 3-(2-sulfonic phenylhydrazo)pentane-2,4-dione (Figure 1a), and 3,7-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane (Figure 1b).

Transition metal–ADB complexes (ADB = azoderivatives of β-diketones) have been successfully applied as homogeneous catalysts for the peroxidative oxidation of alkanes to the corresponding alcohols and ketones, in a few cases with rather high values of TONs.[1,2] In pursuit of this work, the sodium salt of 3-(2-sulfonic phenylhydrazo)pentane-2,4-dione (**SPPD**) was synthesized, fully characterized, and allowed to react with a metal (Cu or Ni) salt in the presence of amines. On the other hand, one of the most important Cage adamantane-like phosphine derivatives of 1,3,5-triaza-7-phosphaadamantane (**PTA**) is 3,7-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane (**DAPTA**), which is prepared by reaction of PTA with acetic anhydride, and possess extremely large molar solubility in water.[3,4] New water soluble copper complex was synthesised by reaction of DAPTA with Cu-salt in mild conditions.

The obtained compounds were characterized by IR spectroscopy, elemental analysis and single crystal X-ray diffraction (**SCXRD**).




Figure 1.

References

- [1] Mahmudov, K. T.; Kopylovich, M. N.; Pombeiro, A. J. L. *Coord. Chem. Rev.* **2013**, *257*, 1244.
- [2] Mahmudov, K. T.; Kopylovich, M. N.; Guedes da Silva, M. F. C.; Pombeiro, A.J.L. *ChemPlusChem* **2014**, *79*, 1523.
- [3] Phillips A. D.; Gonsalvi, L.; Romerosa A.; Vizza, F.; Peruzzini, M. *Coord. Chem. Rev.* **2004**, *248*(11), 955.
- [4] Darendbourg D. J.; Ortiz C. G.; Kamplajn J. W. *Organometallics* **2004**, *23*, 1747.


M–H Bond Repolarization in Hydrogen Bonding and the Proton Transfer Between Two Transition-Metal Hydrides

E.S. Osipova,* O.A. Filippov, V.A. Kirkina, E.I. Gutsul, N.V. Belkova, L.M. Epstein, E.S. Shubina

A. N. Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences, 119991, Vavilov Street 28, Moscow, RUSSIA

Being formally the sources of either hydride ion H⁻ or proton H⁺ neutral transition-metal hydrides demonstrate different reactivity. A computational study of the intermolecular interaction effect on the M–H bond activation in Group 6 metal hydrides ($\eta^5\text{-C}_5\text{H}_5\text{M}(\text{CO})_3\text{H}$ (M = Mo, W) possessing both acidic and basic properties, was carried out. The electronic structure of adducts of hydride complexes ($\eta^5\text{-C}_5\text{H}_5\text{M}(\text{CO})_3\text{H}$ with the Lewis acid BH₃ and base NMe₃ were analysed.¹ It was shown that the nature of orbital interactions in the complex affects the reaction pathway: the $n_{\text{N}} \rightarrow \sigma_{\text{MH}}^*$ interaction leads to deprotonation of the metal hydride and the $\sigma_{\text{MH}} \rightarrow n_{\text{B}}^*$ interaction implies the hydride transfer to the Lewis acid. The ease of M–H bond polarization in complexes ($\eta^5\text{-C}_5\text{H}_5\text{M}(\text{CO})_3\text{H}$ explains their reactivity as donor of either proton or hydride ion.

Previously we reported on the unconventional acid–base interaction between tungsten hydride ($\eta^5\text{-C}_5\text{H}_5\text{W}(\text{CO})_3\text{H}$ which acts as an acid, and nickel(II) pincer hydride (PCP)NiH (PCP = 2,6-(CH₂P(t-C₄H₉)₂)₂C₆H₄) served as a base.² Continuing this line of investigation we studied the proton transfer mechanism between their analogous species bearing hydridotris(pyrazolyl)borate ligand (TpW(CO)₃H) and palladium metal center ((PCP)PdH). Kinetic and thermodynamic parameters of the proton transfer accompanied by dihydrogen evolution were obtained experimentally by means of IR (ν_{MH} , ν_{CO}) and NMR spectroscopy (¹H, ³¹P) in a wide temperature range (190–300 K). The products and intermediates of the process were established and the peculiarities of their structure were revealed by DFT calculations.



Acknowledgements - this work was supported by the RFBR (14-03-00594, 14-03-31074, 15-53-78027)

References

- [1] Filippov O.A., Golub I.E., Osipova E.S., Kirkina V.A., Gutsul E.I., Belkova N.V. *Russ. Chem. Bull.*, **2014**, *11*, 2428 – 2433.
 [2] Levina V.A., Rossin A., Belkova N.V., Chierotti M.R., Epstein L.M., Filippov O.A., Lledós A., Shubina E.S., Zanobini F., Peruzzini M. *Angew. Chem. Int. Ed.*, **2011**, *50*, 1367 –1370.

Catalytic Activity of Copper(I) and Copper(II) 3,5-Dinitro- or 3,5-Bis(Trifluoromethyl)-Pyrazolate Derivatives

Rossana Galassi,^a Oumarou Camille Simon,^{a*} Claudia Graiff,^b M. Fátima C. Guedes da Silva,^c Nuno M. R. Martins,^c Luísa M. D. R. S. Martins,^{c,d} Armando J. L. Pombeiro^c

^a School of Science and Technology, Via S. Agostino 1, 62032 Camerino (MC), Italy.

Email: camillesimon.oumarou@unicam.it

^b Dipartimento di Chimica, Parco Area delle Scienze 17/A, Parma, Italy.

^c Centro de Química Estrutural, Complexo I, Instituto Superior Técnico, Universidade de Lisboa, Portugal.

^d Chemical Engineering Department, Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa, Portugal.

The reaction of [$\mu\text{-N,N-Cu-(3,5}(\text{CF}_3)_2\text{pz})_3$ (**1**) with halide sources led to the formation of Cu(II) hexanuclear derivatives such as [(Cu((CF₃)₂pz)₆(OH)₆)₂ X][Bu₄N] (**2a**)(Cl), (**2b**)(Br), (**2c**)(I), (**2d**)(NO₂).¹ If the reaction occurs in the same conditions but without halides, a highly hydrated hexanuclear metallocycle [(Cu((CF₃)₂pz)₆(OH)₆)₂ (H₂O)_n], (**3**), was isolated, ruling out the template action of the anions. By replacing the CF₃ groups with NO₂ groups in the pyrazole, neither the trinuclear nor the hexanuclear copper derivatives were obtained, and the dinuclear [Cu-(3,5-(NO₂)₂pz)(PPh₃)₂]₂ compound, (**4**), was the only Cu(I) derivative isolated in good yield. Even though CF₃ and NO₂ are both withdrawing groups, a pronounced different chemical behavior was already revealed in the case of similar pyrazolate silver(I) derivatives.² All compounds were characterized by elemental analysis, NMR, IR spectroscopy and ESI-MS spectrometry. In the case of compound **4** its crystal structure has been determined by X Ray diffraction analysis, evidencing its dinuclear nature, build up by two bridging ligands which coordinate two copper atoms through nitrogen donors, forming a six member ring with boat conformation. Compounds (**3**) and (**4**) act as catalysts towards the microwave (MW) assisted peroxidative oxidation of cyclohexane to cyclohexanol and cyclohexanone under mild conditions, which is of industrial significance for the synthesis of Nylon-6,6. High yield (up to 51% of oxygenated products) is obtained after 2h of MW irradiation, using a maximum of 0.2% molar ratio of **3** (the best catalyst) relatively to the substrate in the presence of TEMPO. The efficiency of the catalytic systems as well as the influence of various parameters, such as the reaction time, amount of catalyst, temperature and presence of different additives, are discussed. The work was partially supported by the Fundação para a Ciência e a Tecnologia (project UID/QUI/00100/2013)

References

- [1] Galassi, R.; Burini, A.; Ahmed, A. M. *Eur. J. Inorg. Chem.* **2012**, 3257.
 [2] Galassi, R.; Ricci, S.; Burini, A.; Macchioni A., Rocchigiani, L., Marmottini, F., Tekarli, S. M., Nesterov, V. N. Omary M. A. *Inorg. Chem.*, **2013**, *52*, 14124.

Poster 85

Synthesis of polyboronic acids based on heteroaryl core – synthons for preparation of microporous materials

Krzysztof Gontarczyk,^{*} Krzysztof Durka, Sergusz Luliński

Warsaw University of Technology, Faculty of Chemistry, Laboratory of Fine Technologies
 Noakowskiego 3, 00-664 Warsaw, Poland.
 Email: krzyksiek@gmail.com

Boronic acids derivatives are extensively utilized in modern organic synthesis, especially for carbon-carbon and carbon-heteroatom formation via catalytic coupling.¹ Polyboronic derivatives consisting of more than one boronic group are interesting synthons in material chemistry for preparation of ordered networks.² Such materials – e.g. Covalent Organic Frameworks (COF)³ - have potential applications as adsorbents for gas storage and separation. We have focused on synthesis of di-, tri- and tetraboronic acids⁴ and respective boronate esters based on thienylsilane, bithienyl, bifuryl and 2,4,6-triphenyl-1,3,5-triazine core (Figure 1). Obtained products will be utilized in synthesis of microporous materials. Preliminary results for these materials will also be included.




Figure 1.

References

- [1] Lennox A. J. J., Lloyd-Jones G. C. *Chem. Soc. Rev.* **2014**, *43*, 412.
- [2] Fournier J-H, Maris T., Wuest J.D., Guo W., Galoppini E. *J. Am. Chem. Soc.* **2003**, *125*, 1003.
- [3] Cote, A.; Benin, A. Ockwig, N.; O'Keefe, M.; Matzger, A.; Yaghi, O. *Science* **2005**, *310*, 1166.
- [4] Gontarczyk K., Durka K., Klimkowski P., Luliński S., Serwatowski J., Woźniak K. *J. Organomet. Chem.* **2015**, *783*, 1-9.

Index of Participants

| | Surname | Name | Email | Poster |
|-----|---------------|-----------------|-----------------------------------|--------|
| 1. | Abdalghani | Issam | esam.ghani1988@hotmail.com | 23 |
| 2. | Agonigi | Gabriele | g.ago2@hotmail.it | |
| 3. | Arrigoni | Federica | f.arrigoni5@campus.unimib.it | 15 |
| 4. | Baldino | Salvatore | salvatore.baldino@uniud.it | 51 |
| 5. | Balestri | Davide | davide.balestri@studenti.unipr.it | 13 |
| 6. | Barbieri | Alessia | alessia.barbieri@uniroma1.it | 32 |
| 7. | Barwinski | Bernhard | barwinski@itmc.rwth-aachen.de | |
| 8. | Basalov | Ivan | basalov.vania@yandex.ru | 3 |
| 9. | Battistin | Federica | federica.battistin@phd.units.it | 75 |
| 10. | Biancalana | Lorenzo | lorenzo.biancalana@for.unipi.it | 25 |
| 11. | Boggio | Paolo | paolo.boggio89@gmail.com | 26-64 |
| 12. | Borelli | Tea | tea_borelli@yahoo.it | 18 |
| 13. | Borys | Krzysztof M. | krzysztof.borys@gmail.com | 59 |
| 14. | Bravin | Carlo | carlo.bravin@studenti.unipd.it | 65 |
| 15. | Brusa | Filippo | filippo.brusa@uninsubria.it | 71 |
| 16. | Bucci | Alberto | albechm@gmail.com | 54 |
| 17. | Buccolini | Giulia | giulia.buccolini@gmail.com | |
| 18. | Camille Simon | Oumarou | camillesimon.oumarou@unicam.it | 84 |
| 19. | Capacci | Chiara | chiara.capacci@hotmail.com | 74 |
| 20. | Carminati | Daniela | daniela.carminati@unimi.it | 70 |
| 21. | Cecotka | Piotr | piotrcecotka@gmail.com | |
| 22. | Cesari | Cristiana | cristiana.cesari2@unibo.it | 68 |
| 23. | Chirila | Andrei | a.chirila@uva.nl | 14 |
| 24. | Ciasca | Cosimo Vincenzo | cosimo.ciasca@uniba.it | 50 |
| 25. | Cicco | Luciana | luciana.cicco@uniba.it | 28 |
| 26. | Cingolani | Andrea | andrea.cingolani6@unibo.it | 67 |
| 27. | Condello | Francesca | francesca.condello@unicam.it | 39 |
| 28. | Corbucci | Ilaria | ilariacorbucci@gmail.com | 54 |
| 29. | Coverdale | James | j.p.c.coverdale@warwick.ac.uk | 22 |

| | Surname | Name | Email | Poster |
|-----|----------------|----------------|---------------------------------|---------------|
| 30. | Cybularczyk | Martyna | m_cybularczyk@wp.pl | 20 |
| 31. | Dalla Cort | Antonella | antonella.dallacort@uniroma1.it | |
| 32. | Damas | Liliana | lilianadamas90@gmail.com | 31 |
| 33. | De Nisi | Assunta | assunta.denisi2@unibo.it | |
| 34. | Di Mauro | Giovanni | dm.giovanni91@gmail.com | |
| 35. | Duarte | Tiago | t.duarte@ua.pt | 58 |
| 36. | Esposito | Antonio | anto.exp90@gmail.com | |
| 37. | Esposito | Roberto | roberto.esposito@unina.it | 63 |
| 38. | Fiorini | Valentina | fiorini.valentina@hotmail.it | 8 |
| 39. | Formenti | Dario | dario.formenti@unimi.it | 66 |
| 40. | Giorgi | Simone | simone.giorgi@unicam.it | |
| 41. | Girelli | Chiara Roberta | chiara.girelli@unisalento.it | 12 |
| 42. | Gontarczyk | Krzysztof | krzksiek@gmail.com | 85 |
| 43. | Gorska | Agnieszka | agorska@ch.pw.edu.pl | 55 |
| 44. | Grillo | Giorgio | giorgio_gg@live.com | |
| 45. | Grotenhuis | Colet te | C.teGrotenhuis@uva.nl | 9 |
| 46. | Grudzien | Krzysztof | kgrudzien88@gmail.com | 42 |
| 47. | Jans | Anne | a.c.h.jans@uva.nl | 35 |
| 48. | Jongkind | Lukas | l.j.jongkind@uva.nl | 24 |
| 49. | Jürgens | Eva | eva.juergens@uni-tuebingen.de | 40-78 |
| 50. | Kissel | Alexander | okishvegan@mail.ru | 4 |
| 51. | Kuijpers | Petrus | P.F.Kuijpers@uva.nl | 30 |
| 52. | Licandro | Emanuela | emanuela.licandro@unimi.it | |
| 53. | Lupacchini | Massimiliano | glamo1989@gmail.com | 1 |
| 54. | Macchioni | Alceo | alceo.macchioni@uni.pg | |
| 55. | Maciel | Dina | dmaciel@uma.pt | 47-81 |
| 56. | Mahmoud | Abdallah | abdallah1126@gmail.com | 82 |
| 57. | Malecki | Pawel | pawelmalecki.chem@gmail.com | 53 |
| 58. | Maner | Asif | asif.maner@unicam.it | 57 |
| 59. | Marinelli | Marika | akiramaut@yahoo.it | 41-80 |

| | Surname | Name | Email | Poster |
|-----|------------------|---------------|--|---------------|
| 60. | Martins | Nuno | nunomamartins@tecnico.ulisboa.pt | 38-52 |
| 61. | Mascitti | Andrea | mascio88@gmail.com | 79 |
| 62. | Matos | Cristina | cristina.matos@tecnico.ulisboa.pt | 46 |
| 63. | Matteucci | Elia | elia.matteucci2@unibo.it | 10 |
| 64. | Mazza | Alberto | alberto.mazza@unimi.it | 77 |
| 65. | Michalska | Malina | malina.michalska@gmail.com | 34 |
| 66. | Mirabella | Stefania | stefania.mirabella@unifi.it | 17 |
| 67. | Monari | Magda | magda.monari@unibo.it | |
| 68. | Monticelli | Marco | marco.monticelli.1@studenti.unipd.it | 5 |
| 69. | Nencini | Luca | luca.nencini@unito.it | 49 |
| 70. | Nurttila | Sandra | s.s.nurttila@uva.nl | 27 |
| 71. | Osipova | Elena | aosipova92@gmail.com | 83 |
| 72. | Palmese | Marco | mpalmese42@gmail.com | 6 |
| 73. | Palmucci | Jessica | jessica.palmucci@unicam.it | 19 |
| 74. | Papa | Veronica | veronicapapa@hjournal.it | |
| 75. | Pasqualini | Marta | marta.pasqualini@unicam.it | |
| 76. | Payen | Ricky | ricky.payen@uclouvain.be | 73 |
| 77. | Petrini | Agnese | agnese.petrini@unicam.it | 43 |
| 78. | Ragaini | Fabio | fabio.ragaini@unimi.it | |
| 79. | Renzi | Marco | marco.renzi@unicam.it | |
| 80. | Richard | Sarah | sara.richard@etu.parisdescartes.fr | 7 |
| 81. | Ruddlesden | Amy | ar622@york.ac.uk | 21 |
| 82. | Ruffo | Francesco | ruffo@unina.it | |
| 83. | Sabbatini | Alessandra | alessandra.sabbatini@unicam.it | |
| 84. | Sampaolesi | Susanna | susanna.sampaolesi@unicam.it | |
| 85. | Carmo dos Santos | Nadia | nadiaalessandra.carmodossantos@studenti.unipd.it | 37 |
| 86. | Serrano-Ruiz | Manuel | mserrano@iccom.cnr.it | |
| 87. | Shviro | Meital | meitalshviro@gmail.com | |
| 88. | Silarska | Ewelina | ewelina.silarska@chem.uni.wroc.pl | 33 |
| 89. | Silva Ansaloni | Lilian Marcia | lilian.silva@unimi.it | 11 |

| | Surname | Name | Email | Poster |
|------|----------------|-------------|---------------------------------------|---------------|
| 90. | Singh | Kuldeep | kdsingh@iit.ac.in | 44 |
| 91. | Savourey | Solene | solene.savourey@cea.fr | 62 |
| 92. | Stringer | Tameryn | strtam001@myuct.ac.za | 36 |
| 93. | Szczepaniak | Grzegorz | szczepaniak.grzegorz@gmail.com | 56 |
| 94. | Tarnowicz | Stanislawa | stanislawa.tarnowicz@chem.uni.wroc.pl | 60 |
| 95. | Thakare | Pramod | pramod.thakare@unimi.it | 29 |
| 96. | Tiago | Gonçalo | goncalotiago78@gmail.com | 76 |
| 97. | Tseberlidis | Giorgio | giorgio.tseberlidis@unimi.it | 72 |
| 98. | Von Wolff | Niklas | niklas.von-wolff@cea.fr | 69 |
| 99. | Wirwis | Anna | anna.wirwis@chem.uni.wroc.pl | 2 |
| 100. | Wojcik | Przemyslaw | przemyslaw.wojcik@chem.uni.wroc.pl | 16 |
| 101. | Zaccaria | Francesco | francesco.zaccaria@unina.it | 48 |
| 102. | Zielinski | Grzegorz | gzzielinski@icho.edu.pl | 61 |

